


Physical and Chemical Data on Sediments Deposited in the Missouri and the Mississippi River Flood Plains During the July Through August 1993 Flood

Floods in the Upper Mississippi River Basin 1993

Circular 1120-L



Availability of Publications of the U.S. Geological Survey

Order U.S. Geological Survey (USGS) publications from the offices listed below. Detailed ordering instructions, along with prices of the last offerings, are given in the current-year issues of the catalog "New Publications of the U.S. Geological Survey."

Books, Maps, and Other Publications

By Mail

Books, maps, and other publications are available by mail from—

USGS Information Services
Box 25286, Federal Center
Denver, CO 80225

Publications include Professional Papers, Bulletins, Water-Supply Papers, Techniques of Water-Resources Investigations, Circulars, Fact Sheets, publications of general interest, single copies of permanent USGS catalogs, and topographic and thematic maps.

Over the Counter

Books, maps, and other publications of the U.S. Geological Survey are available over the counter at the following USGS Earth Science Information Centers (ESIC's), all of which are authorized agents of the Superintendent of Documents:

- Anchorage, Alaska—Rm. 101, 4230 University Dr.
- Denver, Colorado—Bldg. 810, Federal Center
- Menlo Park, California—Rm. 3128, Bldg. 3,
345 Middlefield Rd.
- Reston, Virginia—Rm. 1C402, USGS National Center,
12201 Sunrise Valley Dr.
- Salt Lake City, Utah—2222 West, 2300 South (books and
maps available for inspection only)
- Spokane, Washington—Rm. 135, U.S. Post Office
Building, 904 West Riverside Ave.
- Washington, D.C.—Rm. 2650, Main Interior Bldg.,
18th and C Sts., NW.

Maps only may be purchased over the counter at the following USGS office:

- Rolla, Missouri—1400 Independence Rd.

Electronically

Some USGS publications, including the catalog "New Publications of the U.S. Geological Survey" are also available electronically on the USGS's World Wide Web home page at <http://www.usgs.gov>

Preliminary Determination of Epicenters

Subscriptions to the periodical "Preliminary Determination of Epicenters" can be obtained only from the Superintendent of

Documents. Check or money order must be payable to the Superintendent of Documents. Order by mail from—

Superintendent of Documents
Government Printing Office
Washington, DC 20402

Information Periodicals

Many Information Periodicals products are available through the systems or formats listed below:

Printed Products

Printed copies of the Minerals Yearbook and the Mineral Commodity Summaries can be ordered from the Superintendent of Documents, Government Printing Office (address above). Printed copies of Metal Industry Indicators and Mineral Industry Surveys can be ordered from the Center for Disease Control and Prevention, National Institute for Occupational Safety and Health, Pittsburgh Research Center, P.O. Box 18070, Pittsburgh, PA 15236-0070.

Mines FaxBack: Return fax service

1. Use the touch-tone handset attached to your fax machine's telephone jack. (ISDN [digital] telephones cannot be used with fax machines.)
2. Dial (703) 648-4999.
3. Listen to the menu options and punch in the number of your selection, using the touch-tone telephone.
4. After completing your selection, press the start button on your fax machine.

CD-ROM

A disc containing chapters of the Minerals Yearbook (1993–95), the Mineral Commodity Summaries (1995–97), a statistical compendium (1970–90), and other publications is updated three times a year and sold by the Superintendent of Documents, Government Printing Office (address above).

World Wide Web

Minerals information is available electronically at <http://minerals.er.usgs.gov/minerals/>

Subscription to the catalog "New Publications of the U.S. Geological Survey"

Those wishing to be placed on a free subscription list for the catalog "New Publications of the U.S. Geological Survey" should write to—

U.S. Geological Survey
903 National Center
Reston, VA 20192

**PHYSICAL AND CHEMICAL DATA ON SEDIMENTS
DEPOSITED IN THE MISSOURI AND THE MISSISSIPPI
RIVER FLOOD PLAINS DURING THE JULY THROUGH
AUGUST 1993 FLOOD**

By Gregg K. Schalk, Robert R. Holmes, Jr., and Gary P. Johnson

Floods in the Upper Mississippi River Basin, 1993

U.S. GEOLOGICAL SURVEY CIRCULAR 1120-L

U.S. DEPARTMENT OF THE INTERIOR
BRUCE BABBITT, Secretary

U.S. GEOLOGICAL SURVEY
Charles G. Groat, Director

Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government

1998

Library of Congress Cataloging In Publication Data

Schalk, Gregg K., 1968—

Physical and chemical data on sediments deposited in the Missouri and the Mississippi River flood plains during the July through August 1993 flood / by Gregg K. Schalk, Robert R. Holmes, Jr., and Gary P. Johnson.

p. 62 cm. — (Floods in the upper Mississippi River basin, 1993; L) (U.S. Geological Survey circular ; 1120)

Includes bibliographical references. Supt. of Docs. no. : I 19.4/2:1120-L ISBN 0-607-88279-4

1. River sediments—Mississippi River Valley. 2. River sediments—Missouri River Valley. 3. Floods—Mississippi River Valley.

4. Floods—Missouri River Valley. I. Holmes, Robert R., 1965— . II. Johnson, Gary P., 1969— . III. Title. IV. Series

V. Series: U.S. Geological Survey circular ; 1120.

GB1217.F58 1993 Ch. L [GB1399.8] 551.48'9'0977 s--dc21 [551.3'54'0977]

97-49113

CIP

Free on application to the
U.S. Geological Survey
Branch of Information Services
Box 25286, Federal Center
Denver, CO 80225

FOREWORD

During spring and summer 1993, record flooding inundated much of the upper Mississippi River Basin. The magnitude of the damages—in terms of property, disrupted business, and personal trauma—was unmatched by any other flood disaster in United States history. Property damage alone is expected to exceed \$10 billion. Damaged highways and submerged roads disrupted overland transportation throughout the flooded region. The Mississippi and the Missouri Rivers were closed to navigation before, during, and after the flooding. Millions of acres of productive farmland remained under water for weeks during the growing season. Rills and gullies in many tilled fields are the result of the severe erosion that occurred throughout the Midwestern United States farmbelt. The hydrologic effects of extended rainfall throughout the upper Midwestern United States were severe and widespread. The banks and channels of many rivers were severely eroded, and sediment was deposited over large areas of the basin's flood plain. Record flows submerged many areas that had not been affected by previous floods. Industrial and agricultural areas were inundated, which caused concern about the transport and fate of industrial chemicals, sewage effluent, and agricultural chemicals in the floodwaters. The extent and duration of the flooding caused numerous levees to fail. One failed levee on the Raccoon River in Des Moines, Iowa, led to flooding of the city's water treatment plant. As a result, the city was without drinking water for 19 days.

As the Nation's principal water-science agency, the U.S. Geological Survey (USGS) is in a unique position to provide an immediate assessment of some of the hydrological effects of the 1993 flood. The USGS maintains a hydrologic data network and conducts extensive water-resources investigations nationwide. Long-term data from this network and information on local and regional hydrology provide the basis for identifying and documenting the effects of the flooding. During the flood, the USGS provided continuous streamflow and related information to the National Weather Service (NWS), the U.S. Army Corps of Engineers, the Federal Emergency Management Agency (FEMA), and many State and local agencies as part of its role to provide basic information on the Nation's surface- and ground-water resources at thousands of locations across the United States. The NWS has used the data in forecasting floods and issuing flood warnings. The data have been used by the Corps of Engineers to operate water diversions, dams, locks, and levees. The FEMA and many State and local emergency management agencies have used USGS hydrologic data and NWS forecasts as part of the basis of their local flood-response activities. In addition, USGS hydrologists are conducting a series of investigations to document the effects of the flooding and to improve understanding of the related processes. The major initial findings from these studies will be reported in this Circular series as results become available.

U.S. Geological Survey Circular 1120, *Floods in the Upper Mississippi River Basin, 1993*, consists of individually published chapters that will document the effects of the 1993 flooding. The series includes data and findings on the magnitude and frequency of peak discharges; precipitation; water-quality characteristics, including nutrients and man-made contaminants; transport of sediment; assessment of sediment deposited on flood plains; effects of inundation on ground-water quality; flood-discharge volume; effects of reservoir storage on flood peaks; stream-channel scour at selected bridges; extent of floodplain inundation; and documentation of geomorphologic changes.



Charles G. Groat
Dirctor

CONTENTS

Foreword.....	III
Abstract.....	1
Introduction	1
Purpose and Scope.....	1
Description of Study Area	2
Study Methods	3
Site Selection and Location	4
Sampling Techniques.....	5
Laboratory Analytical Methods.....	6
Quality-Assurance Samples	11
Physical and Chemical Data on Sediments.....	12
Particle-Size Distribution.....	12
Moisture Content and Volatile Solids, Nutrient, and Carbon Concentrations.....	12
Selected Element Concentrations	12
Pesticide Concentrations.....	12
Semivolatile Organic Compound Concentrations	21
Summary and Conclusions	21
References Cited	21

FIGURES

1. Map showing location of study area and sediment deposition sites sampled during September 1993 in the Missouri and the Mississippi River flood plains.....	2
2.-4. Photographs showing:	
2. Silt and clay sediment deposited in the Missouri River flood plain near Hermann, Missouri, during the 1993 flood	3
3. Sand dunes deposited in the Missouri River flood plain near St. Charles, Missouri, during the 1993 flood.....	3
4. Aerial photograph showing the Missouri River flood plain near Hartsburg, Missouri, November 1993.	3

TABLES

1. Selected data for sampling sites in the Missouri and the Mississippi River flood plains	5
2. Detection or reporting limits of analytical methods used for the determination of constituent concentrations in the less than 2-millimeter size fraction of postflood sediment samples.....	6
3. Detection limits and methods used for the determination of element concentrations in the less than 62-millimeter size fraction of postflood sediment samples	8
4. Detection limits used for the determination of pesticide and semivolatile organic compound concentrations in the less than 2-millimeter size fraction of postflood sediment samples.....	10
5. Particle-size distribution of postflood sediment and preflood soil samples in the Missouri and the Mississippi River flood plains.....	13
6. Moisture content and volatile solids, nutrient, and carbon concentrations in the less than 2-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains	16
7. Concentrations of elements and carbon in the less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains.....	24
8. Descriptive statistics of selected element and carbon concentrations in the less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains.....	35
9. Measured and expected concentrations of elements in standard reference material	18
10. Estimated pesticide concentrations in the less than 2-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains	19
11. Concentrations of semivolatile organic compounds in the less than 2-millimeter size fraction of postflood sediment samples in the Missouri River flood plain, sample set reagent spike, standard reference material, and surrogate samples	46
12. Concentrations of semivolatile organic compounds in the less than 2-millimeter size fraction of postflood sediment samples in the Mississippi River flood plain, sample set reagent spike, standard reference material, and surrogate samples	50
13. Tentatively identified compounds that were detected in the gas chromatography/mass spectrometry analysis.....	53

CONVERSION FACTORS AND VERTICAL DATUM

Multiply	By	To obtain
millimeter (mm)	0.03937	inch
meter (m)	3.281	foot
square meter (m^2)	10.76	square foot
kilometer (km)	0.6214	mile
kilogram (kg)	2.205	pound
gram (g)	0.03527	ounce
square kilometer (km^2)	0.3861	square mile

Temperature in degrees Celsius ($^{\circ}\text{C}$) can be converted to degrees Fahrenheit ($^{\circ}\text{F}$) as follows:

$$\text{Celsius temperature } (^{\circ}\text{C}) \qquad ^{\circ}\text{F} = 1.8^{\circ}\text{C} + 32 \qquad \text{Fahrenheit temperature } (^{\circ}\text{F})$$

Sea level: In this report, “sea level” refers to the National Geodetic Vertical Datum of 1929 (NGVD of 1929)—a geodetic datum derived from a general adjustment of the first-order level nets of both the United States and Canada, formerly called Sea Level Datum of 1929.

Physical and Chemical Data on Sediments Deposited in the Missouri and the Mississippi River Flood Plains During the July through August 1993 Flood

By Gregg K. Schalk, Robert R. Holmes, Jr., and Gary P. Johnson

Abstract

In July and August 1993, Missouri and Mississippi River floodwaters overtopped or breached levees and inundated agricultural and urban areas in the flood plains. To assess the regional variability of quality of silt and clay sediments transported and deposited during inundation, sediment samples were collected in September 1993 at 25 sites in the Missouri and the Mississippi River flood plains. The samples were analyzed for physical properties and chemical constituents that included particle-size distribution, water content, volatile solids, nutrients, carbon, selected trace elements, pesticides, and semivolatile organic compounds for postflood sediment samples, and particle-size distribution for preflood soil samples. Pesticides were detected in samples from many of the sites, with alachlor detected in one sample at a concentration of 54 micrograms per kilogram. Many semivolatile organic compounds were detected in samples at several sites, with the concentration of fluoranthene in one sample at 270 milligrams per kilogram. Procedures for selecting sites, techniques developed for sampling, laboratory analytical methods, and quality-assurance sampling methods also are described.

INTRODUCTION

Peak discharges exceeded 100-year recurrence intervals at U.S. Geological Survey (USGS) streamflow-gaging stations on the Missouri and the Mississippi Rivers during July and August 1993 (Parrett and

others, 1993). Floodwaters inundated the flood plain as a result of failed levees. Sediments transported by the floodwaters were deposited in varying thicknesses in these inundated areas. The deposited silt and clay sediments were thought to contain elevated concentrations of nutrients and trace elements. Also, because the floodwaters inundated various agricultural and urban areas, several pesticides and organic compounds were thought to have attached to silt and clay sediments that were mobilized and deposited downstream in the inundated flood plain. Large concentrations of these various constituents are of concern because of their potential effects on croplands and wildlife habitat in the flood plain. To obtain information that would be needed to address these concerns, the USGS conducted a study to assess the regional variability in the physical and chemical characteristics of silt- and clay-size (hereafter referred to as "silt and clay") sediments deposited in the Missouri and the Mississippi River flood plains. The study included determination of the presence of contaminants in the sediments and a comparison of particle-size distribution in postflood sediment and pre-flood soil at several locations.

Purpose and Scope

This report describes the sampling and analytical techniques, quality-assurance methods, and the results of analyses of sediment samples collected in September 1993 from 25 sites along the flood plains of the Missouri and the Mississippi Rivers. Data presented for the postflood sediment samples include particle-size distribution, water content, volatile solids, and concentrations of nutrients, carbon, other selected elements, pesticides, and semivolatile organic compounds. The particle-size distribution of preflood soil sampled beneath the postflood sediment samples also is presented.

Description of Study Area

The study area was the flood plains of the Missouri and the Mississippi Rivers that were inundated by floodwaters during the July through August 1993

flood—a 560-river mile reach of the Missouri River from Nebraska City, Nebraska, to St. Louis, Missouri, and a 578-river mile reach of the Mississippi River from Dubuque, Iowa, to Cairo, Illinois (fig. 1). These flood plains contain a series of levees constructed to




Figure 1. Location of study area and sediment deposition sites sampled during September 1993 in the Missouri and the Mississippi River flood plains.

withstand floodwaters from the rivers and their tributaries. The predominant land use in the flood plain is agriculture, interspersed with urban areas. Postflood sediment samples were collected along the flood plains at 25 sites.

Deposition of silt and clay sediment in the flood plain generally was associated with "slack water" areas where floodwaters entered with little velocity and pooled. These areas commonly were located near the flood-plain side of the levee adjacent to the levee break where the flow paths separated. Silt and clay deposition was determined to be as much as 140 mm deep in the flood plain of the Missouri River near Hermann, Missouri (fig. 2).

Although sites with sand deposition generally were not sampled, sand dunes as deep as 4 m were observed near many levee breaks, such as those near St. Charles, Missouri (fig. 3). Sand dunes along the flood plain near Hartsburg, Missouri, were as long as 3,000 m and as high as 3 m (fig. 4).

STUDY METHODS

For this study, 25 sites were selected to sample postflood sediment and preflood soil. Site selection and location techniques, sampling techniques, laboratory analytical methods, and quality-assurance samples are discussed in the following sections.



Figure 2. Silt and clay sediment deposited in the Missouri River flood plain near Hermann, Missouri, during the 1993 flood. The photograph was taken on September 7, 1993.




Figure 3. Sand dunes deposited in the Missouri River flood plain near St. Charles, Missouri, during the 1993 flood. The photograph was taken on September 7, 1993.



Figure 4. Aerial photograph showing the Missouri River flood plain near Hartsburg, Missouri, November 1993. The river length is about 20 kilometers.

Site Selection and Location

Twelve sites along the Missouri River flood plain and 13 sites along the Mississippi River flood plain, in areas where substantial quantities of silt and clay were deposited during the 1993 flood, were selected for sampling. Approximate locations for the sampling sites initially were selected from topographic maps. The sites were located about 45 river miles apart—those on the Missouri River extending upstream from the junction of the Missouri and the Mississippi Rivers, and the sites along the Mississippi River beginning at Dubuque, Iowa, and extending to the junction of the Mississippi and the Ohio Rivers (fig. 1; table 1).

Exact locations for sampling were chosen on the basis of onsite reconnaissance, which was limited to 5 river miles upstream or downstream from the initially selected approximate location. Factors considered in selection of the sampling location included the degree of disturbance, abundance of silt and clay, accessibility of the location, and the type of preflood land use. The preferred land use was agricultural production fields. Areas of possible contamination as a result of human or animal activities were avoided. After visual inspection of the surface of the silt and clay deposit, a hole was dug to permit inspection of the structure of the deposit and the total deposition depth (fig. 2). A deposition depth of at least 0.2 m was required before a sample was collected.

Table 1. Selected data for sampling sites in the Missouri and the Mississippi River flood plains

[MO, Missouri; MS, Mississippi]

Site no. (fig. 1)	U.S. Geological Survey station number	Latitude/longitude, in degrees, minutes, seconds ¹	Date sampled	Elevation, in feet	River mile ²	Depth interval for postflood sediment samples, in millimeters ³
Missouri River flood-plain samples						
MO.1	384125090341401	384125.4 0903414.2	09-07-93	451	38	0-70
MO.2	384305091263301	384304.5 0912632.9	09-07-93	505	98	20-140
MO.3	383438092034301	383438.4 0920343.2	09-08-93	537	137	0-70
MO.4	385142092265701	385141.6 0922656.9	09-08-93	570	176	0-30
	385142092265702	385141.6 0922656.9	09-08-93	570	176	30-150
MO.5	391823092565401	391823.2 0925654.2	09-09-93	620	238	0-40
MO.6	391714093225301	391713.9 0932252.9	09-09-93	648	281	0-30
MO.7	391152094100401	391152.2 0941003.6	09-10-93	710	338	0-60
MO.8	391134094432401	391134.0 0944324.2	09-10-93	748	380	0-40
MO.9	393857095015401	393857.1 0950154.1	09-10-93	790	432	0-90
MO.10	395247095074001	395247.1 0950740.1	09-11-93	836	474	0-90
MO.11	401901095370101	401900.9 0953700.6	09-11-93	883	525	0-80
MO.12	404010095485601	404010.0 0954855.7	09-11-93	919	560	0-20

Table 1. Selected data for sampling sites in the Missouri and the Mississippi River flood plains—Continued

[MO, Missouri; MS, Mississippi]

Site no. (fig. 1)	U.S. Geological Survey station number	Latitude/longitude, in degrees, minutes, seconds ¹	Date sampled	Elevation, in feet	River mile ²	Depth interval for postflood sediment samples, in millimeters ³
Mississippi River flood-plain samples						
MS.1	422840090381801	422839.7 0903818.2	09–18–93	597	578	0–100
MS.2	420450090112601	420449.9 0901126.0	09–18–93	585	536	0–40
MS.3	413026090363201	413026.3 0903631.8	09–18–93	550	481	0–60
MS.4	411023091022301	411022.8 0910222.7	09–19–93	537	435	0–40
MS.5	403851091142501	403851.0 0911424.6	09–20–93	530	389	0–70
MS.6	400642091252201	400641.5 0912521.5	09–20–93	483	341	0–20
MS.7	393318091073101	393318.2 0910731.1	09–21–93	458	291	0–60
MS.8	390239090435901	390239.3 0904359.0	09–21–93	440	245	0–20
MS.9	385224090123801	385223.9 0901237.9	09–22–93	423	203	0–60
MS.10	381248090165601	381248.1 0901656.2	09–22–93	395	148	0–20
MS.11	374532089404501	374532.0 0894044.9	09–24–93	365	95	0–110
MS.12	371745089302201	371744.9 0893021.9	09–23–93	335	52	0–60
MS.13	370333089135201	370332.8 0891352.4	09–23–93	315	10	0–20

¹The first number listed is latitude; the second number is longitude.²For the Missouri River, 0 river mile is at the junction of the Missouri and the Mississippi Rivers; for the Mississippi River, 0 river mile is at the junction of the Mississippi and the Ohio Rivers.³0 millimeter is the surface of the sediment deposition.

Once the exact sampling site was selected, a hand-held global positioning system (GPS), accurate to within 30 m, was used to determine the latitude and longitude of the site. Latitude, longitude, USGS station number, description, date sampled, elevation of the site, the location in river miles with respect to either the Missouri River (MO prefix site numbers) or the Mississippi River (MS prefix site numbers), and the sediment-deposition depth are given in table 1. Elevations were determined to ± 3 m from 7.5-minute USGS topographic maps. River miles were determined to ± 0.5 km from 7.5-minute USGS topographic maps.

Sampling Techniques

To collect a postflood sediment sample, a hole was dug to identify the interface between the postflood deposition and the preflood soil. The flood-deposited sediments were well consolidated, whereas the underlying, preflood soil typically had a “loose” structure and commonly contained roots and other organic debris. After digging the hole, the surface area was cleared of any organic debris. The vertical cut area was shaved with a clean plastic scraper to ensure no contamination. The plastic scraper, a plastic scoop, and latex gloves were used to collect a sample from the vertical cut. The sample was collected above the interface

in a 1-m² area or less, depending on the quantity of sediment available. A scoop of the sediment was collected at various points in the sample area. The sediment was composited, placed in a sample container, and immediately chilled. After sample collection, the plastic scrapers and scoops were cleaned and rinsed with deionized water, methanol, and organic-free water. No samples were collected where water had not receded from the area. One postflood sediment sample was collected at all sites except site MO.4; at this site, two samples mostly composed of sand were collected—one from 0 to 30 mm and the other from 30 to 150 mm.

Samples collected for moisture and carbon content and for concentrations of volatile solids, nutrients, pesticides, and semivolatile organic compounds were analyzed only in the less than 2-mm size fraction. Because the postflood sediment samples were determined onsite to have a particle diameter of less than 2 mm, no samples were sieved.

After the postflood sediment samples were collected, preflood soil samples were collected for particle-size distribution. All flood deposition was removed from the sample area with a clean plastic scraper to the postflood deposition and preflood soil interface. Preflood soil samples were collected by using the same collection procedures used for the postflood sediment samples.

Different sample containers were used for each analysis type. Samples for analysis of particle-size distribution were collected in a plastic bag and sealed. Samples for analysis of moisture content, volatile solids, nutrients, and carbon were collected in a wide-mouth polyethylene bottle with a plastic lid. Samples

for analysis of pesticides and semivolatile organic compounds were collected in a 1-L wide-mouth glass bottle with a Teflon-lined lid that had been baked at 450° C. Samples for analysis of other selected elements were collected in a wide-mouth plastic bottle with a Teflon-lined lid.

Laboratory Analytical Methods

Samples for particle-size distribution were analyzed at the USGS laboratory in Rolla, Missouri. The following classification of particle-size diameter (Guy, 1969) was used to describe material in this report: gravel, 2 to less than 64 mm; sand, 0.062 to less than 2 mm; silt, 0.04 to less than 0.062 mm; and clay, less than 0.004 mm. Silt and clay material are considered to be fine grained; sand- and gravel-size (hereafter referred to as "sand and gravel") material are considered to be coarse. Samples that contained sand, silt, and clay material were analyzed by using a hydrometer, visual accumulation (VA) tube, and wet sieve (Guy, 1969). Samples that contained mostly silt and clay material were analyzed by using a hydrometer as described by the U.S. Army Corps of Engineers (1970).

Samples for analysis of water content and concentrations of volatile solids, nutrients, and carbon in the less than 2-mm size fraction were analyzed at the USGS laboratory in Arvada, Colorado, by using analytical methods having detection or reporting limits listed in table 2.

Table 2. Detection or reporting limits of analytical methods used for the determination of constituent concentrations in the less than 2-millimeter size fraction of postflood sediment samples

[Analysis done at the U.S. Geological Survey laboratory, Arvada, Colorado]

Constituent	Detection or reporting limit	Constituent	Detection or reporting limit
Moisture content by dry weight, in percent..	0.1	Solids, volatile, in milligrams per kilogram.....	1.0
Nutrients, in milligrams per kilogram			
Nitrite plus nitrate as nitrogen	2.0	Ammonia plus organic nitrogen.....	20
Ammonia as nitrogen2	Phosphorus	40
Carbon, in grams per kilogram			
Total	0.1	Inorganic.....	0.1
Pesticides, in micrograms per kilogram¹			
Alachlor	3.6	Dieldrin	3.2
Atrazine	6.8	Metolachlor	3.6
4,4'-DDE	4.0	Simazine	3.2
Desethylatrazine	1.2	Trifluralin	4.8

Table 2. Detection or reporting limits of analytical methods used for the determination of constituent concentrations in the less than 2-millimeter size fraction of postflood sediment samples—Continued

[Analysis done at the U.S. Geological Survey laboratory, Arvada, Colorado]

Constituent	Detection or reporting limit	Constituent	Detection or reporting limit
Semivolatile organic compounds, in micrograms per kilogram			
Acenaphthene.....	10	p-Cresol.....	10
Acenaphthylene.....	5.0	Dibenzo[<i>a,h</i>]anthracene.....	20
Acridine.....	10	Dibenzothiphene	5.0
Anthracene	5.0	Di- <i>n</i> -butylphthalate	10
Anthraquinone.....	20	1,2-Dichlorobenzene.....	10
Azo-benzene.....	20	1,3-Dichlorobenzene.....	10
Benzo[<i>a</i>]anthracene	10	1,4-Dichlorobenzene.....	10
Benzo[<i>b</i>]fluoranthene.....	10	2,4-Dichlorophenol.....	20
Benzo[<i>k</i>]fluoranthene	10	Diethyl phthalate.....	10
Benzo[<i>g,h,i</i>]perylene	20	1,2-Dimethylnaphthalene.....	10
Benzo[<i>a</i>]pyrene	10	1,6-Dimethylnaphthalene.....	10
Benzo[<i>c</i>]quinoline.....	10	2,6-Dimethylnaphthalene.....	10
2,2'-Biquinoline	20	3,5-Dimethylphenol.....	20
4-Bromophenyl phenyl ether.....	10	Dimethyl phthalate.....	10
Butylbenzylphthalate	10	4,6-Dinitro-2-methylphenol.....	200
C ₈ -Alkyl-phenol.....	10	2,4-Dinitrophenol.....	300
9H-Carbazole	10	2,4-Dinitrotoluene.....	20
bis(2-Chloroethoxy)methane	10	2,6-Dinitrotoluene.....	50
bis(2-Chloroethyl)ether.....	10	di- <i>n</i> -Octyl phthalate.....	20
bis(2-Chloroisopropyl)ether.....	20	bis(2-Ethylhexyl)phthalate.....	10
4-Chloro-3-methylphenol.....	30	2-Ethynaphthalene	10
2-Chloronaphthalene	10	Fluoranthene	5.0
2-Chlorophenol	10	9H-Fluorene	10
4-Chlorophenyl phenyl ether.....	10	Hexachlorobenzene	10
Chrysene.....	10	Hexachlorobutadiene	20
Hexachlorocyclopentadiene	20	<i>n</i> -Nitrosodiphenylamine	10
Hexachloroethane.....	20	Pentachloroanisol.....	20
Indeno[<i>1,2,3-cd</i>]pyrene	20	Pentachloronitrobenzene	30
Isophorone.....	10	Pentachlorophenol	100
Isoquinoline.....	10	Phenanthrene.....	5.0
1-Methyl-9H-fluorene	10	Phenanthridine	10
2-Methylanthracene	10	Phenol	10
4,5-Methylenephенanthrene.....	20	Pyrene	5.0
1-Methylphenanthrene	10	Quinoline	10
1-Methylpyrene	10	2,3,5,6-Tetramethylphenol	30
Naphthalene	10	1,2,4-Trichlorobenzene	10
Nitrobenzene	10	2,4,6-Trichlorophenol	20
2-Nitrophenol	100	2,3,6-Trimethylnaphthalene	10
4-Nitrophenol	40	2,4,6-Trimethylphenol	30
<i>n</i> -Nitroso di- <i>n</i> -propylamine	10		

¹Reporting limits are estimates (E.T. Furlong, U.S. Geological Survey, written commun., 1993).

Water content and volatile solids concentration were determined by using methods described in Fishman and Friedman (1989). Nutrient samples were analyzed by using methods described in Fishman (1993). Total and inorganic carbon samples were analyzed by using methods described in Wershaw and others (1987). Organic carbon concentrations were computed by subtracting the inorganic carbon from the total carbon concentrations.

The concentrations of selected elements and carbon in the less than 0.062-mm size fraction were determined at the USGS laboratory in Lakewood, Colorado, by using the methods described in Arbogast (1990). The constituents analyzed and descriptions of the different methods of analysis for each constituent are listed in table 3. The total carbon concentration was determined by using methods described in Wershaw and others (1987).

Table 3. Detection limits and methods used for the determination of element concentrations in the less than 62-millimeter size fraction of postflood sediment samples

[4N HCl, 4 normal hydrochloride acid; mg/kg, milligrams per kilogram; 0.1N HCl, 0.1 normal hydrochloric acid; ICP-AES, inductively coupled plasma-atomic emission spectrometry; IR, infrared spectrometry; n.a., not analyzed; HGAAS, hydride-generation atomic absorption spectrometry; FAAS, flame atomic absorption spectrometry; CVAAS, cold-vapor atomic absorption spectrometry; DN, delayed neutron activation—methods presented in Arbogast (1990). Analyzed at the U.S. Geological Survey laboratory, Lakewood, Colorado]

Element	Method	Total digestion ¹	Detection limit	
			4N HCl digestion ² (mg/kg)	0.1N HCl digestion ³ (mg/kg)
Aluminum	ICP-AES	0.005 percent by weight	1.3	1.3
Calcium	—do—	.005 percent by weight	1.0	1.0
Iron	—do—	.02 percent by weight	.40	.40
Magnesium	—do—	.005 percent by weight	5.8	5.8
Phosphorus	—do—	.005 percent by weight	12.5	12.5
Potassium	—do—	.01 percent by weight	7.5	7.5
Sodium	—do—	.006 percent by weight	2.5	2.5
Sulfur	IR ⁴	.05 percent by weight	n.a.	n.a.
Titanium	ICP-AES	.005 percent by weight	.25	.25
Antimony	HGAAS	0.2 mg/kg	n.a.	n.a.
	ICP-AES	n.a.	6.3	6.3
Arsenic	—do—	10 mg/kg	3.8	3.8
	HGAAS	.2 mg/kg	.05	.05
Barium	ICP-AES	1 mg/kg	.10	.10
Beryllium	—do—	1 mg/kg	.025	.025
Bismuth	—do—	10 mg/kg	7.5	7.5
Cadmium	—do—	2 mg/kg	.50	.50
	FAAS ⁵	.1 mg/kg	n.a.	n.a.
Cerium	ICP-AES	5 mg/kg	5.0	5.0
Chromium	—do—	2 mg/kg	1.5	1.5
Cobalt	—do—	2 mg/kg	1.3	1.3
Copper	—do—	2 mg/kg	.25	.25
Europium	—do—	2 mg/kg	n.a.	n.a.
Gallium	—do—	4 mg/kg	n.a.	n.a.
Gold	—do—	8 mg/kg	n.a.	n.a.
Holmium	—do—	4 mg/kg	n.a.	n.a.
Lanthanum	—do—	2 mg/kg	.75	.75
Lead	—do—	4 mg/kg	6.3	6.3
Lithium	—do—	2 mg/kg	.38	.38
Manganese	—do—	4 mg/kg	.63	.63
Mercury	CVAAS ⁴	.02 mg/kg	n.a.	.006
Molybdenum	ICP-AES	2 mg/kg	1.0	1.0

Table 3. Detection limits and methods used for the determination of element concentrations in the less than 62-millimeter size fraction of postflood sediment samples—Continued

[4N HCl, 4 normal hydrochloride acid; mg/kg, milligrams per kilogram; 0.1N HCl, 0.1 normal hydrochloric acid; ICP-AES, inductively coupled plasma-atomic emission spectrometry; IR, infrared spectrometry; n.a., not analyzed; HGAAS, hydride-generation atomic absorption spectrometry; FAAS, flame atomic absorption spectrometry; CVAAS, cold-vapor atomic absorption spectrometry; DN, delayed neutron activation—methods presented in Arbogast (1990). Analyzed at the U.S. Geological Survey laboratory, Lakewood, Colorado]

Element	Method	Total digestion ¹	Detection limit	
			4N HCl digestion ² (mg/kg)	0.1N HCl digestion ³ (mg/kg)
Neodymium	—do—	9 mg/kg	n.a.	n.a.
Nickel	—do—	3 mg/kg	1.3	1.3
Niobium	—do—	4 mg/kg	n.a.	n.a.
Scandium	—do—	2 mg/kg	n.a.	n.a.
Selenium	HGAAS	.2 mg/kg	.05	.05
Silver.....	ICP-AES FAAS ⁵	2 mg/kg .1 mg/kg	0.50 n.a.	0.50 n.a.
Strontium.....	ICP-AES	2 mg/kg	.025	.025
Tantalum.....	—do—	40 mg/kg	n.a.	n.a.
Thorium.....	—do—	6 mg/kg	n.a.	n.a.
.....	DN	1 mg/kg	n.a.	n.a.
Tin.....	ICP-AES	5 mg/kg	10	10
Tungsten	—do—	n.a.	7.5	7.5
Uranium.....	—do—	100 mg/kg	n.a.	n.a.
.....	DN	.2 mg/kg	n.a.	n.a.
Vanadium.....	ICP-AES	2 mg/kg	.38	.38
Ytterbium.....	—do—	1 mg/kg	n.a.	n.a.
Yttrium	—do—	2 mg/kg	.25	.25
Zinc.....	—do—	2 mg/kg	.25	.25
Zirconium	—do—	n.a.	.63	.63
Carbon, total	Combustion ⁴	.05 percent by weight	n.a.	n.a.
Carbon, organic	Difference ⁴	.05 percent by weight	n.a.	n.a.
Carbon, carbonate as CO ₂	Titration ⁴	.01 percent by weight	n.a.	n.a.

¹Concentrated hydrochloric, hydrofluoric, nitric, and perchloric acids.

²Conditionally available fraction in the sediments.

³Available fraction in the sediments.

⁴Method presented in Arbogast (1990).

⁵Method presented in O'Leary and Viets (1986).

Selected pesticide and semivolatile organic compound concentrations were determined for all sites in the less than 2-mm size fraction at the Enseco Analytical Laboratory, Arvada, Colorado. A list of constituents and the detection limits is given in table 4. Samples for pesticides were analyzed by using U.S. Environmental Protection Agency (USEPA) method SW8080, and samples for semivolatile organic compounds were analyzed by using USEPA method SW8270 (U.S. Environmental Protection Agency, 1986).

Concentrations of selected pesticides and semivolatile organic compounds were determined for 15 of 25 sites in the less than 2-mm size fraction at the USGS laboratory in Arvada. A list of constituents and the detection limits is given in table 2. The detection limits for 4,4'-DDE (hereafter referred to in this report as "DDE") and dieldrin were lower than the limits used by the Enseco Analytical Laboratory. A different procedure was used by the USGS to analyze for the pesticides and semivolatile organic compounds.

Table 4. Detection limits used for the determination of pesticide and semivolatile organic compound concentrations in the less than 2-millimeter size fraction of postflood sediment samples

[All detection limits in milligrams per kilogram. Analyzed at the Enseco Analytical Laboratory, Arvada, Colorado]

Compound	Detection limit	Compound	Detection limit
Pesticides¹			
Aldrin	0.010	Endosulfan II	0.020
alpha-BHC010	Endosulfan sulfate020
beta-BHC010	Heptachlor010
delta-BHC010	Heptachlor epoxide010
gamma-BHC (Lindane)010	Methoxychlor	
Chlordane10	PCB, Aroclor 101610
4,4'-DDD020	PCB, Aroclor 122110
4,4'-DDE020	PCB, Aroclor 123210
4,4'-DDT020	PCB, Aroclor 124210
Dieldrin020	PCB, Aroclor 124810
Endrin020	PCB, Aroclor 125410
Endrin aldehyde020	PCB, Aroclor 126010
Endosulfan I010	Toxaphene20
Semivolatile organic compounds²			
Acenaphthene	0.50	Chrysene	0.50
Acenaphthylene50	Dibenzofuran50
Acetophenone	2.5	Dibenzo[<i>a,h</i>]anthracene50
4-Aminobiphenyl	2.5	Di- <i>n</i> -butylphthalate50
Aniline	2.5	1,2-Dichlorobenzene50
Anthracene50	1,3-Dichlorobenzene50
Benzidine	5.6	1,4-Dichlorobenzene50
Benzo[<i>a</i>]anthracene50	2,4-Dichlorophenol50
Benzo[<i>b</i>]fluoranthene50	2,6-Dichlorophenol	2.5
Benzo[<i>k</i>]fluoranthene50	Diethyl phthalate50
Benzoic acid	2.5	<i>p</i> -Dimethylaminoazobenzene	2.5
Benzo[<i>g,h,i</i>]perylene50	7,12-Dimethylbenzo[<i>a</i>]anthracene	2.5
Benzo[<i>a</i>]pyrene50	2,4-Dimethylphenol50
Benzyl alcohol	1.0	Dimethyl phthalate50
4-Bromophenyl phenyl ether50	4,6-Dinitro-2-methylphenol	1.5
Butylbenzylphthalate50	2,4-Dinitrophenol	1.5
4-Chloroaniline	1.0	2,4-Dinitrotoluene50
bis(2-Chloroethoxy)methane50	2,6-Dinitrotoluene50
bis(2-Chlorooethyl)ether50	Di- <i>n</i> -octyl phthalate50
bis(2-Chloroisopropyl)ether50	Diphenylamine	2.5
4-Chloro-3-methylphenol50	bis(2-Ethylhexyl)phthalate50
2-Chloronaphthalene50	Ethyl methanesulfonate	2.5
2-Chlorophenol50	Fluoranthene50
4-Chlorophenyl phenyl ether50	Fluorene50
1-Chloronaphthalene	2.5	Hexachlorobenzene50

Table 4. Detection limits used for the determination of pesticide and semivolatile organic compound concentrations in the less than 2-millimeter size fraction of postflood sediment samples—Continued

[All detection limits in milligrams per kilogram. Analyzed at the Enseco Analytical Laboratory, Arvada, Colorado]

Compound	Detection limit	Compound	Detection limit
Semivolatile organic compounds ² —Continued			
Hexachlorobutadiene	0.50	4-Nitrophenol	2.5
Hexachlorocyclopentadiene50	<i>n</i> -Nitrosodimethylamine	2.5
Hexachloroethane50	<i>n</i> -Nitrosodi- <i>n</i> -butylamine	2.5
Indeno[1,2,3- <i>cd</i>]pyrene50	<i>n</i> -Nitrosodi- <i>n</i> -propylamine50
Isophorone50	<i>n</i> -Nitrosodiphenylamine50
3-Methylcholanthrene	2.5	<i>n</i> -Nitrosodiperidine	2.5
Methyl methanesulfonate.....	2.5	Pentachlorobenzene	2.5
2-Methylnaphthalene50	Pentachloronitrobenzene	2.5
2-Methylphenol.....	.50	Pentachlorophenol	1.5
4-Methylphenol.....	.50	Phenacetin	2.5
Naphthalene50	Phenanthrene50
1-Naphthylamine.....	2.5	Phenol50
2-Naphthylamine.....	2.5	Pyrene	2.5
2-Nitroaniline	2.5	1,2,4,5-Tetrachlorobenzene	2.5
3-Nitroaniline	2.5	2,3,4,6-Tetrachlorophenol	2.5
4-Nitroaniline	2.5	1,2,4-Trichlorobenzene50
Nitrobenzene50	2,4,5-Trichlorophenol	1.5
2-Nitrophenol50	2,4,6-Trichlorophenol50

¹Method SW8080 (U.S. Environmental Protection Agency, 1986).

²Method SW8270 (U.S. Environmental Protection Agency, 1986).

Quality-Assurance Samples

Quality-assurance samples were collected and analyzed to determine the variability in sampling and analytical techniques. Replicate samples were collected at sites MO.7, MS.1, MS.7, and MS.13 to determine the variability of sampling techniques. Duplicate samples were split from samples collected for analysis of concentrations of semivolatile organic compounds and selected elements to determine the variability in analytical techniques.

Sample set reagent spikes were used to determine the accuracy of the analyses for pesticides and semivolatile organic compounds. During the gel permeation chromatography (GPC) process, pentachlorophenol (chlorophenols) may have been sorbed and desorbed, which caused low and variable recoveries, and degradation of hexachlorocyclopentadiene and hexachlorobutadiene likely occurred in the injection port of the gas chromatography/mass spectrometry (GC/MS) (E.T. Furlong, U.S. Geological Survey, writ-

ten commun., 1993) and affected the concentrations detected in the reagent spikes.

The effect of the sample matrix and gross processing errors in samples collected for pesticides and semivolatile organic compounds can be monitored by the results of the surrogate compound recoveries. Selected surrogate compounds with chemical properties similar to some of the analytes of interest are added to each sample at the beginning of extraction.

Standard reference materials (SRM) for elements, pesticides, and semivolatile organic compounds were analyzed to determine laboratory analytical accuracies. Analytical split analyses for the elemental SRM also were performed to determine precision of the analyses.

Analysis of sample set blanks was performed for samples collected for pesticides and semivolatile organic compounds. Analysis of a blank determines if any interferences occurred during the analytical process of the sample set. No interferences were detected in any of the blanks for either sample set.

PHYSICAL AND CHEMICAL DATA ON SEDIMENTS

The following sections present and summarize the results of analyses of particle-size distribution, water content, and the concentrations of nutrients, volatile solids, carbon, other selected elements, pesticides, and semivolatile organic compounds in postflood sediment samples and the particle-size distribution of pre-flood soil samples.

The mean and the standard deviation were computed for water content, volatile solids, nutrients, carbon, selected elements, and pesticides. The mean and the standard deviations were computed only for constituents detected in more than 50 percent of the samples. In those samples where the constituent concentration was less than the detection limit, one-half the detection limit was used to compute the mean and the standard deviation.

Particle-Size Distribution

The particle-size distribution data for the postflood sediment and preflood soil samples are presented in table 5. Because of the way the sample sites were selected, the postflood sediment samples primarily are silt and clay except for the deeper sample from MO.4, which contained 91.7 percent by weight sand. Postflood sediment samples that were greater than 50 percent by weight clay were from sites MO.3, MO.6, MO.8, MO.9, and MS.9 (fig. 1) and ranged from 52.3 to 78.2 percent by weight clay. All other postflood sediment samples ranged from 51.5 to 87.7 percent by weight silt. The sample from MO.1 had the lowest clay content of 8.0 percent by weight. Sand in postflood sediment samples ranged from 0 percent by weight at MO.6 to 28.4 percent by weight at MS.13.

Samples of preflood soil were collected from areas of various land uses, which included agricultural production fields, vacant urban areas, vacant land, and levees. The preflood soil sample collected from the base of a levee at MS.5 had the highest gravel content—7.5 percent by weight. Sand in preflood soil samples ranged from less than 1 to 82.5 percent by weight, silt content ranged from 9.0 to 85.2 percent by weight, and clay content ranged from 1.0 to 71.5 percent by weight.

Moisture Content and Volatile Solids, Nutrient, and Carbon Concentrations

The data for moisture content and for concentrations of volatile solids, nutrients, and carbon for all samples in the less than 2-mm size fraction are presented in table 6. Moisture content in the postflood sediment ranged from 7 to 51 percent by weight. Volatile solids concentrations ranged from 6,750 to 91,500 mg/kg. The deeper sample from MO.4 had the smallest nutrient concentrations detected in all samples. The nitrite plus nitrate as nitrogen concentration of 140 mg/kg in the MS.12 sample was about three times larger than the concentrations from other sites. For all samples except MO.4 layer 2, total carbon was present predominantly as organic carbon. The total carbon concentration from this sample (2.5 g/kg) was an order of magnitude less than the concentration in other samples.

Selected Element Concentrations

Generally, as grain size decreases in bottom-sediment material, element concentrations increase (Horowitz, 1991, p. 16). On the basis of this general correlation, postflood sediment samples were analyzed for concentrations of 12 selected elements in the silt and clay fraction. These data are listed in table 7, descriptive statistics are listed in table 8 (both follow “References Cited” section), and the results for SRM for elements are in listed table 9.

Pesticide Concentrations

Samples were analyzed for pesticides by two laboratories. The method and detection limit for the pesticides for samples analyzed at the Enseco Analytical Laboratory are listed in table 4 and at the USGS laboratory in Arvada are listed in table 2. No pesticides were detected by the Enseco Analytical Laboratory. Results of pesticide analyses by the USGS laboratory are listed in table 10. Some of the concentrations reported in table 10 are less than the detection or reporting limit listed in table 2 because pesticide concentrations were determined by a custom-USGS method that generally has lower detection limits. Only DDE and dieldrin were analyzed by both laboratories.

Aalachlor was detected in 14 samples; the highest concentration of 54 µg/kg was in the MS.5 sample.

Table 5. Particle-size distribution of postflood sediment and preflood soil samples in the Missouri and the Mississippi River flood plains

[MO, Missouri; MS, Mississippi; —, no data. Analyzed at the U.S. Geological Survey sediment laboratory, Rolla, Missouri]

Site no. (fig. 1)	Sample type	Gravel						Sand						Silt						Clay	
		8.0	4.0	2.0	1.0	0.50	0.25	0.125	0.062	0.031	0.016	0.008	0.004	0.002	0.001	0.0005	0.0002	0.00005	0.00000	Clay	
Missouri River																					
MO.1	Postflood.....	100	100	100	100	100	99.4	99.3	98.2	89.4	60.4	30.7	12.3	8.0	5.0	—	—	—	—		
	Preflood.....	100	100	100	100	100	100	100	87.9	45.0	42.9	27.0	20.0	15.2	—	—	—	—	—		
MO.2	Postflood.....	100	100	100	100	100	100	100	99.4	95.2	88.4	70.4	51.1	39.6	27.3	—	—	—	—		
	Preflood.....	100	100	100	100	100	100	99.6	99.2	97.0	90.1	81.6	64.2	49.2	37.4	—	—	—	—		
MO.3	Postflood.....	100	100	100	100	100	100	100	99.8	99.4	97.3	79.9	63.3	52.3	38.8	—	—	—	—		
	Preflood.....	100	100	100	100	100	100	98.8	93.3	89.9	87.9	82.1	72.2	60.4	50.6	—	—	—	—		
MO.4	Postflood ¹	100	100	100	100	100	100	100	99.7	99.0	92.4	75.6	31.1	16.6	11.8	11.5	—	—	—		
	Postflood ²	100	99.5	99.2	98.4	93.8	85.6	24.9	7.5	1	—	—	—	—	—	—	—	—	—		
	Preflood.....	96.6	96.1	95.5	93.9	89.5	85.6	76.0	54.1	27.3	16.6	10.3	8.1	—	—	—	—	—	—		
MO.5	Postflood.....	100	100	100	100	100	99.7	99.2	98.9	98.8	75.0	67.7	51.5	38.1	30.8	—	—	—	—		
	Preflood.....	100	100	100	100	100	100	96.9	69.7	36.9	29.1	15.9	10.8	7.9	—	—	—	—	—		
MO.6	Postflood.....	100	100	100	100	100	100	100	100	100	100	99.2	98.1	92.2	78.2	57.2	—	—	—		
	Preflood.....	100	100	100	100	100	100	99.9	99.8	99.7	97.2	95.6	93.3	71.5	53.3	—	—	—	—		
MO.7	Postflood.....	100	100	100	100	100	99.9	99.5	99.0	98.8	94.9	72.6	52.4	39.7	29.1	—	—	—	—		
	Replicate.....	100	100	100	100	100	100	99.9	99.6	98.2	64.0	38.9	26.6	20.2	13.5	—	—	—	—		
	Split.....	100	100	100	100	100	100	100	99.8	97.2	85.2	43.4	30.7	23.9	21.1	—	—	—	—		
	Preflood.....	100	100	100	100	99.6	99.6	98.4	96.5	84.0	38.8	32.6	25.2	16.2	7.7	—	—	—	—		
MO.8	Postflood.....	100	100	100	100	100	100	100	100	99.9	98.3	91.7	78.3	60.8	44.8	—	—	—	—		
	Preflood.....	100	100	100	100	100	100	98.9	97.9	91.8	54.0	43.9	26.2	22.8	—	—	—	—	—		
MO.9	Postflood.....	100	100	100	100	100	100	100	100	99.9	96.7	93.3	83.6	66.3	48.4	—	—	—	—		
	Preflood.....	100	100	100	100	100	100	100	100	100	39.2	23.4	21.4	18.2	15.2	13.8	—	—	—	—	

Table 5. Particle-size distribution of postflood sediment and preflood soil samples in the Missouri and the Mississippi River flood plains—Continued

[MO, Missouri; MS, Mississippi; —, no data. Analyzed at the U.S. Geological Survey sediment laboratory, Rolla, Missouri]

Site no. (fig. 1)	Sample type	Gravel						Sand						Silt						Clay										
		8.0	4.0	2.0	1.0	0.50	0.25	0.125	0.062	0.031	0.016	0.008	0.004	8.0	4.0	2.0	1.0	0.50	0.25	0.125	0.062	0.031	0.016	0.008	0.004	0.002				
Missouri River—Continued																														
MO.10	Postflood.....	100	100	100	100	100	100	100	100	99.8	98.5	86.6	81.3	64.1	45.2	33.1	33.1	33.1	33.1	33.1	33.1	33.1	33.1	33.1	33.1	33.1				
	Preflood.....	100	100	100	100	100	100	100	99.1	97.4	93.6	81.3	17.3	14.0	10.4	3.6	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7			
MO.11	Postflood.....	100	100	100	100	99.9	99.9	99.9	99.8	99.5	87.7	48.8	34.1	24.4	19.1	19.1	19.1	19.1	19.1	19.1	19.1	19.1	19.1	19.1	19.1	19.1	19.1			
	Preflood.....	100	100	100	100	100	100	100	96.6	87.9	71.1	62.0	50.8	45.4	30.4	25.7	25.7	25.7	25.7	25.7	25.7	25.7	25.7	25.7	25.7	25.7	25.7	25.7		
MO.12	Postflood.....	100	100	100	100	100	100	100	100	99.9	99.5	85.4	62.0	42.5	31.3	22.6	22.6	22.6	22.6	22.6	22.6	22.6	22.6	22.6	22.6	22.6	22.6	22.6		
	Preflood.....	100	100	100	100	100	100	99.6	98.4	95.5	91.1	67.7	50.5	39.7	32.1	25.8	25.8	25.8	25.8	25.8	25.8	25.8	25.8	25.8	25.8	25.8	25.8	25.8	25.8	
Mississippi River																														
MS.1	Postflood.....	100	100	100	100	100	100	100	99.8	99.7	99.7	67.6	40.0	21.0	12.0	6.8	6.8	6.8	6.8	6.8	6.8	6.8	6.8	6.8	6.8	6.8	6.8	6.8		
	Replicate.....	100	100	100	100	100	100	100	100	99.6	99.3	76.3	51.5	32.0	24.4	11.5	11.5	11.5	11.5	11.5	11.5	11.5	11.5	11.5	11.5	11.5	11.5	11.5		
	Preflood.....	100	100	100	99.1	99.1	99.1	98.5	98.2	97.7	97.2	72.8	50.3	29.0	18.2	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7	7.7		
MS.2	Postflood.....	100	100	100	100	100	100	99.9	99.9	99.8	99.8	95.2	83.0	63.0	35.4	9.9	9.9	9.9	9.9	9.9	9.9	9.9	9.9	9.9	9.9	9.9	9.9	9.9	9.9	
	Preflood.....	100	100	99.2	98.6	98.6	95.7	93.7	91.8	89.3	46.5	25.7	15.8	15.8	15.1	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8	7.8	
MS.3	Postflood.....	100	100	100	100	99.9	99.9	99.6	99.0	94.6	71.4	36.2	22.2	18.9	12.8	12.8	12.8	12.8	12.8	12.8	12.8	12.8	12.8	12.8	12.8	12.8	12.8	12.8		
	Preflood.....	100	100	100	100	99.6	97.8	96.8	95.7	94.6	71.8	41.2	24.7	24.7	14.3	10.1	10.1	10.1	10.1	10.1	10.1	10.1	10.1	10.1	10.1	10.1	10.1	10.1	10.1	
MS.4	Postflood.....	100	100	100	99.6	92.1	70.9	56.3	53.3	51.2	48.6	44.0	34.3	34.3	23.1	15.6	15.6	15.6	15.6	15.6	15.6	15.6	15.6	15.6	15.6	15.6	15.6	15.6	15.6	
	Preflood.....	100	100	98.6	92.5	85.0	34.6	26.2	19.2	10.0	2.8	1.6	1.2	1.0	.9	.9	.9	.9	.9	.9	.9	.9	.9	.9	.9	.9	.9	.9		
MS.5	Postflood.....	100	100	96.9	92.5	85.0	34.6	26.2	19.2	10.0	2.8	1.6	1.2	1.0	.9	.9	.9	.9	.9	.9	.9	.9	.9	.9	.9	.9	.9	.9		
	Preflood.....	100	100	100	100	100	100	99.9	99.6	99.2	98.8	84.7	60.8	33.0	20.3	20.3	20.3	20.3	20.3	20.3	20.3	20.3	20.3	20.3	20.3	20.3	20.3	20.3		
MS.6	Postflood.....	100	100	99.5	98.7	98.5	98.3	98.1	87.1	79.7	57.3	33.1	33.1	33.1	33.1	28.0	28.0	28.0	28.0	28.0	28.0	28.0	28.0	28.0	28.0	28.0	28.0	28.0	28.0	28.0

Table 5. Particle-size distribution of postflood sediment and preflood soil samples in the Missouri and the Mississippi River flood plains—Continued

[MO, Missouri; MS, Mississippi; —, no data. Analyzed at the U.S. Geological Survey sediment laboratory, Rolla, Missouri]

Site no. (fig. 1)	Sample type	Sand						Silt				Clay	
		8.0	4.0	2.0	1.0	0.50	0.25	0.125	0.062	0.031	0.016	0.008	0.004
Mississippi River—Continued													
MS.7	Postflood.....	100	100	100	100	100	100	99.9	99.5	97.5	52.8	35.5	23.6
	Replicate.....	100	100	100	100	100	99.9	98.9	98.9	78.3	56.6	37.2	17.8
	Preflood.....	100	100	99.0	96.0	83.6	72.4	69.3	68.5	66.7	58.3	45.7	31.1
MS.8	Postflood.....	100	100	100	99.5	99.4	99.2	98.8	98.3	91.4	70.3	50.1	31.4
	Preflood.....	100	100	99.7	98.9	98.7	98.1	97.0	97.0	76.5	55.7	33.8	18.1
MS.9	Postflood.....	100	100	100	100	100	100	100	99.8	98.6	97.9	83.5	62.7
	Preflood.....	100	10	100	100	100	100	99.0	92.5	83.4	73.1	61.7	51.8
MS.10	Postflood.....	100	100	100	100	99.8	99.8	99.7	99.5	58.4	32.2	21.4	18.2
	Preflood.....	100	98.7	98.5	98.5	98.5	97.9	95.8	83.4	71.5	53.3	35.5	26.2
MS.11	Postflood.....	100	100	100	100	100	100	100	98.7	75.2	57.9	42.0	32.6
	Preflood.....	100	100	100	100	100	100	100	99.9	94.6	84.0	70.7	58.6
MS.12	Postflood.....	100	100	100	100	100	100	99.9	99.9	89.7	77.6	60.1	45.1
	Preflood.....	100	100	99.9	99.9	98.9	87.8	51.6	31.3	25.7	22.9	12.1	10.1
MS.13	Postflood.....	100	100	100	100	100	100	100	71.6	46.6	27.2	15.4	9.1
	Replicate.....	100	100	100	100	100	100	100	78.2	66.8	43.2	32.7	20.5
	Preflood.....	100	100	100	100	100	100	99.6	98.9	42.4	30.9	18.0	10.4

¹Layer 1, 0 to 30 millimeters.

²Layer 2, 30 to 150 millimeters.

Table 6. Moisture content and volatile solids, nutrient, and carbon concentrations in the less than 2-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains

[USGS, U.S. Geological Survey, mg/kg, milligrams per kilogram; N, nitrogen, g/kg, grams per kilogram; MO, Missouri; rep, replicate sample; MS, Mississippi; —, no data. All analyses by the USGS laboratory, Arvada, Colorado, except as indicated]

Site no. (fig. 1)	Date	Time	Moisture content, in percent by weight			Carbon				
			USGS labora- tory	Enseco Analytical Laboratory	Solids, volatile (mg/kg)	Nitrite + nitrate as N (mg/kg)	Ammonia as N (mg/kg)	Ammonia + organic nitrogen as N (mg/kg)	Phosphorus, total (mg/kg)	Total (g/kg)
Missouri River										
MO.1	09-07-93	1040	26	29	28,700	9	3.9	800	440	11
MO.2	09-07-93	1636	27	36	9,110	8	7.2	710	390	11
MO.3	09-08-93	1125	30	36	74,600	20	5.2	2,300	760	23
¹ MO.4	09-08-93	1605	37	28	68,400	18	6.3	1,900	760	21
² MO.4	09-08-93	1610	7	8.3	6,750	7	3.2	70	210	25
MO.5	09-09-93	1315	35	23	46,000	41	7.5	1,100	620	51
MO.6	09-09-93	1820	37	26	73,800	33	6.8	2,100	750	21
MO.7	09-10-93	1000	26	24	34,500	16	7.6	1,200	560	13
MO.7 rep	09-10-93	1001	35	28	67,500	16	7.2	2,000	740	21
MO.8	09-10-93	1415	38	32	65,000	23	4.2	2,100	740	18
MO.9	09-10-93	1745	35	30	63,200	13	7.3	1,900	810	19
MO.10	09-11-93	1000	25	23	45,500	26	3.3	1,200	560	13
MO.11	09-11-93	1415	26	33	32,700	30	13	940	560	12
MO.12	09-11-93	1730	30	36	30,600	31	14	800	550	14
Mean			28	—	42,000	20	6.9	1,200	580	17
Standard deviation			8.2	—	22,000	10	3.5	670	180	12
Mississippi River										
MS.1	09-18-93	1110	35	34	48,200	21	5.8	2,200	680	24
MS.1 rep	09-18-93	1111	35	35	52,300	21	5.3	2,000	660	22
MS.2	09-18-93	1720	51	50	85,300	15	13	3,200	990	32
										.8

Table 6. Moisture content and volatile solids, nutrient, and carbon concentrations in the less than 2-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

[USGS, U.S. Geological Survey; mg/kg, milligrams per kilogram; N, nitrogen; g/kg, grams per kilogram; MO, Missouri; rep., replicate sample; MS, Mississippi; —, no data. All analyses by the USGS laboratory, Arvada, Colorado, except as indicated]

Site no. (fig. 1)	Date	Time	Moisture content, in percent by weight			Carbon					
			USGS labora- tory	Enseco Analytical Laboratory	Solids, volatile (mg/kg)	Nitrite + nitrate as N (mg/kg)	Ammonia as N (mg/kg)	Ammonia + organic nitrogen as N (mg/kg)	Phosphorus, total (mg/kg)	Total (g/kg)	Organic (g/kg)
Mississippi River—Continued											
MS.3	09-19-93	1440	34	33	35,400	10	7.1	1,400	640	22	17
MS.4	09-19-93	1845	50	48	91,500	19	8.0	3,300	920	38	34
MS.5	09-20-93	0920	27	35	50,900	29	5.6	1,600	690	18	16
MS.6	09-20-93	1730	48	51	64,900	15	60	2,600	790	25	25
MS.7	09-21-93	0940	48	42	77,600	14	8.9	2,500	680	28	25
MS.7 rep	09-21-93	0941	48	51	71,500	11	24	2,400	680	29	26
MS.8	09-21-93	1545	26	28	43,000	47	9.7	1,400	310	15	14
MS.9	09-22-93	0830	32	33	50,900	44	6.2	1,300	610	14	12
MS.10	09-22-93	1420	26	25	51,300	42	7.4	1,400	270	15	15
MS.11	09-24-93	0920	36	34	75,400	70	15	2,100	740	23	22
MS.12	09-23-93	1120	31	33	71,700	140	24	1,900	850	21	20
MS.13	09-23-93	1525	28	28	27,500	39	7.8	1,700	460	10	8.3
MS.13 rep	09-23-93	1526	27	28	23,900	12	6.1	1,700	390	10	8.0
Mean			36	—	60,000	39	14	2,000	660	22	20
Standard deviation			9.6	—	20,000	35	15	680	220	78	7.4
Missouri and Mississippi Rivers											
Mean			32	—	51,000	30	10	1,700	620	20	18
Standard deviation			9.6	—	22,000	28	11	770	200	10	10

¹Layer 1, 0 to 30 millimeters.

²Layer 2, 30 to 150 millimeters.

Table 9. Measured and expected concentrations of elements in standard reference material

[GXR-2, U.S. Geological Survey (USGS) geochemical exploration sample; GSD-6, geochemical reference sample from China; STD, standard deviation; wt. %, percent by weight; mg/kg, milligrams per kilogram; —, no data; (), concentration does not meet criteria for recommended concentrations; L, detected at less than the lower reporting limit but not quantified; AAS, atomic absorption spectrometry. All elements analyzed by inductively coupled plasma-atomic emission spectrometry unless otherwise noted. Analyzed at the USGS laboratory, Lakewood, Colorado]

Element	Unit	GXR-2 concentrations				GSD-6 concentrations			
		Split sample				Split sample			
		Measured	Measured	Expected ¹	STD ¹	Measured	Measured	Expected ²	STD ²
Aluminum.....	wt. %	6.3	6.5	18.57	0.06	7.6	7.6	7.49	—
Calcium.....	wt. %	.91	.92	.88	.04	2.8	2.8	2.77	—
Iron.....	wt. %	1.9	1.9	1.84	.05	4.1	4.1	(4.12)	—
Magnesium.....	wt. %	.79	.80	.88	.02	1.8	1.8	1.81	—
Phosphorus.....	wt. %	.07	.07	.068	—	.11	.11	.10	0.07
Potassium.....	wt. %	1.4	1.4	1.36	.04	2.0	2.0	2.03	—
Sodium.....	wt. %	.59	.58	.54	.08	1.7	1.7	1.71	—
Sulfur, total.....	wt. %	.05L	.05L	.0315	—	.09	.10	(.08)	—
Titanium.....	wt. %	.25	.25	.29	.01	.40	.39	.46	.02
Antimony.....	mg/kg	40	38	47	4	1.6	1.8	1.25	.33
Arsenic.....	mg/kg	24	23	21	4	15	16	13.6	1.5
Arsenic, AAS.....	mg/kg	24	23	21	4	15	15	13.6	1.5
Barium.....	mg/kg	2,200	2,200	2,210	160	310	310	330	40
Beryllium.....	mg/kg	1.0	1.0	1.6	—	1.0	1.0	1.7	.4
Bismuth.....	mg/kg	10.0L	10.0L	.44	.1	10.0L	10.0L	5	.6
Cadmium.....	mg/kg	4.0	4.0	4.1	.04	2.0L	2.0L	.43	.04
Cadmium, AAS.....	mg/kg	4.5	4.2	4.1	.04	.35	.39	.43	.04
Cerium.....	mg/kg	50	50	50	—	70	67	68	8
Chromium.....	mg/kg	35	34	34	4	200	200	190	24
Cobalt.....	mg/kg	10	9.0	9.0	1.3	29	28	24.4	3
Copper.....	mg/kg	82	84	70	6	410	410	383	18
Europium.....	mg/kg	2.0L	2.0L	.77	—	2.0L	2.0L	1.5	.13
Gallium.....	mg/kg	19	18	37	—	18	16	16.7	.8
Gold.....	mg/kg	8.0L	8.0L	.046	1.019	8.0L	8.0L	—	—
Holmium.....	mg/kg	4.0L	4.0L	—	—	4.0L	4.0L	(.78)	.17
Lanthanum.....	mg/kg	27	29	25	3	38	37	.79	8
Lead.....	mg/kg	590	590	670	50	24	25	.27	5
Lithium.....	mg/kg	64	63	48	—	44	44	.40	2
Manganese.....	mg/kg	1,000	1,000	980	40	970	970	970	60
Mercury.....	mg/kg	2.9	2.9	2.7	.6	.02	.02	.045	.008
Molybdenum.....	mg/kg	2.0L	2.0L	1.5	.2	5.0	6.0	.77	1.2
Neodymium.....	mg/kg	21	22	—	—	32	30	.73	6
Nickel.....	mg/kg	19	19	20	4	84	83	.78	7
Niobium.....	mg/kg	10	11	9.0	—	12	11	12	4.6
Scandium.....	mg/kg	6.0	6.0	7.2	—	16	16	17	1.6
Selenium.....	mg/kg	.7	.7	.57	.17	.3	.3	(.3)	.13
Silver.....	mg/kg	17	16	18	3	2.0L	2.0L	.36	.04
Silver, AAS.....	mg/kg	19	19	18	3	.38	.36	.36	.04
Strontium.....	mg/kg	160	160	152	16	280	280	266	28
Tantalum.....	mg/kg	40.0L	40.0L	.78	—	40.0L	40.0L	.72	.07
Thorium.....	mg/kg	9.0	9.0	8.6	.2	9.0	8.0	9.0	2
Tin.....	mg/kg	5.0L	5.0L	1.2	.4	5.0L	5.0L	2.8	1
Uranium.....	mg/kg	100.0L	100.0L	1.98	1.25	100.0L	100.0L	2.4	.5
Vanadium.....	mg/kg	51	50	61	9	140	140	142	12
Ytterbium.....	mg/kg	1.0	1.0	2	—	2.0	2.0	2.1	.4
Yttrium.....	mg/kg	15	15	19	—	20	20	20.2	2.4
Zinc.....	mg/kg	550	550	460	30	140	140	144	10
Carbon, total.....	wt. %	2.9	2.8	2.5	.24	.93	.84	(1.17)	—
Carbon, organic....	wt. %	2.9	2.8	—	—	.40	.33	(.62)	—
Carbon, carbonate..	wt. %	.01L	.01	—	—	.53	.51	(.55)	—
Carbon dioxide.....	wt. %	.02	.04	—	—	1.9	1.9	—	—

¹Gladney and others (1984).²Xie and Yan (1985).

Table 10. Estimated pesticide concentrations in the less than 2-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains

[Concentrations in micrograms per kilogram. DDE, dichlorodiphenyl dichloroethylene; MO, Missouri; <, less than; —, no data; dup., laboratory duplicate sample; rep., field replicate sample; blank, blank sample; n.a., not applicable; MS, Mississippi. Analyzed at the U.S. Geological Survey laboratory, Arvada, Colorado]

Site no. (fig. 1)	Missouri River						Mississippi River					
	A1achlor	Atrazine	4,4'-DDE	Desethyl- atrazine	Dieldrin	Metolachlor	Simazine	Trifluralin	A1achlor	Atrazine	4,4'-DDE	Desethyl- atrazine
MO.1	2.0	4.2	<4.0	<1.2	6.2	3.2	—	5.1	—	—	—	—
MO.2	6.9	6.0	.65	<1.2	7.1	5.1	—	5.5	—	—	—	—
¹ MO.4	.80	7.8	.77	<1.2	4.4	3.3	—	4.8	—	—	—	—
² MO.4 dup	<3.6	<6.8	<4.0	<1.2	<3.2	<3.6	—	<4.8	—	—	—	—
MO.5	<3.6	<6.8	<4.0	<1.2	<3.2	<3.6	—	<4.8	—	—	—	—
	3.6	6.0	1.6	<1.2	3.9	4.2	—	5.1	—	—	—	—
MO.7	4.6	6.0	2.5	<1.2	5.6	4.4	—	6.0	—	—	—	—
MO.7 rep	3.8	4.6	3.6	<1.2	5.2	3.5	—	7.6	—	—	—	—
MO.8	18	10	.64	2.9	3.6	9.9	—	4.8	—	—	—	—
MO.10	2.6	7.6	.36	<1.2	2.5	6.8	—	<4.8	—	—	—	—
MO.11	12	9.5	.87	<1.2	<3.2	5.6	—	5.4	—	—	—	—
MO blank	<3.6	<6.8	<4.0	<1.2	<3.2	<3.6	—	<4.8	—	—	—	—
Mean	6.3	7.1	1.2	n.a.	4.8	5.3	n.a.	4.9	—	—	—	—
Standard deviation	5.9	2.0	.76	n.a.	1.6	2.2	n.a.	1.1	—	—	—	—
Mississippi River												
MS.1	9.5	<6.8	0.93	<1.2	<3.2	3.4	6.4	<4.8	—	—	—	—
MS.1 rep	11	<6.8	<4.0	<1.2	<3.2	4.1	6.4	<4.8	—	—	—	—
MS.3	40	<6.8	1.1	<1.2	<3.2	6.1	<3.2	<4.8	—	—	—	—
MS.5	54	9.1	.96	<1.2	<3.2	7.5	<3.2	4.9	—	—	—	—
MS.7	27	6.7	.94	<1.2	<3.2	4.3	<3.2	<4.8	—	—	—	—
MS.7 rep	22	6.8	1.2	<1.2	<3.2	4.2	<3.2	<4.8	—	—	—	—

Table 10. Estimated pesticide concentrations in the less than 2-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

[Concentrations in micrograms per kilogram. DDE, dichlorodiphenyl dichloroethylene; MO, Missouri; <, less than; —, no data; dup, laboratory duplicate sample; rep, field replicate sample; blank, blank sample; n.a., not applicable; MS, Mississippi. Analyzed at the U.S. Geological Survey laboratory, Arvada, Colorado]

Site no. (fig. 1)	Aalachlor	Atrazine	4,4'-DDE	Desethyl- atrazine	Dieldrin	Metolachlor	Simazine	Trifluralin
Mississippi River—Continued								
MS.9	4.2	9.1	0.32	<1.2	<3.2	4.1	<3.2	5.9
MS.11	3.2	5.1	<4.0	<1.2	<3.2	2.9	<3.2	<4.8
MS.11 dup	2.9	4.6	.69	<1.2	<3.2	2.8	<3.2	4.9
MS.13	<3.6	4.2	.48	<1.2	<3.2	1.9	<3.2	18
MS.13 rep	<3.6	4.2	.47	<1.2	<3.2	1.9	<3.2	13
MS blank	<3.6	<6.8	<4.0	<1.2	<3.2	<3.6	<3.2	<4.8
Mean	20	5.8	.96	n.a.	n.a.	4.3	n.a.	n.a.
Standard deviation	21	2.5	.54	n.a.	n.a.	1.9	n.a.	n.a.
Missouri and Mississippi Rivers								
Mean	13	6.5	1.1	n.a.	n.a.	4.8	n.a.	n.a.
Standard deviation	16	2.2	.66	n.a.	n.a.	2.1	n.a.	n.a.
Standard reference material								
Measured	<3.6	<6.8	9.1	<1.2	<3.2	<3.6	<3.2	<4.8
Expected ³	—	—	9.71	—	—	—	—	—

¹Layer 1, 0 to 30 millimeters.

²Layer 2, 30 to 150 millimeters.

³Noncertified concentration.

Atrazine and DDE were detected in 13 samples. Desethylatrazine was detected only in one sample. Dieldrin was detected in seven samples, all of which were from the Missouri River flood-plain sites. Metolachlor was detected in 15 samples. The Missouri River flood-plain site samples were not analyzed for simazine, and only one Mississippi River site (MS.1) had a concentration that was larger than the detection or reporting limit. Trifluralin was detected in 10 samples.

Semivolatile Organic Compound Concentrations

Sediment samples were analyzed for semivolatile organic compounds by two laboratories. None of these compounds were detected by the Enseco Analytical Laboratory, which used USEPA analytical methods. However, the compounds were detected in the same samples by the USGS laboratory in Arvada, which used analytical methods that have lower detection limits than the methods used by Enseco. Eight semivolatile organic compounds were detected in sediment samples from eight sites in the Missouri River flood plain (table 11, which follows the "References Cited" section). Twenty-five semivolatile organic compounds were detected in samples from 7 sites in the Mississippi River flood plain (table 12, which follows the "References Cited" section); concentrations of the compounds detected in these samples ranged from 23 µg/kg of 2,6-dimethylnaphthalene at MS.5 and MS.11 and 2-ethylnaphthalene at MS.3 to 270 µg/kg of fluoranthene at MS.3. The number of compounds detected per site ranged from 5 to 24.

Identification of unknown organic compounds that were detected in the GC/MS analysis is presented in table 13, which follows the "References Cited" section. The siloxane compound group is present in silicon-based grease in the sampling or laboratory equipment, but is not present in the sediment. In the Missouri River flood-plain samples, 11 compounds and 4 compound groups were identified, and in the Mississippi River flood-plain samples, 29 compounds and 7 compound groups were identified.

SUMMARY AND CONCLUSIONS

Floodwaters inundated the Missouri and Mississippi River flood plains during July and August 1993 as a result of peak discharges that exceeded 100-year

recurrence intervals at streamflow-gaging stations. Sediments transported by the floodwaters were deposited in varying thicknesses in the inundated areas. Because the sediments were thought to contain elevated concentrations of nutrients and trace elements, the U.S. Geological Survey collected sediment samples at 25 sites in the Missouri and the Mississippi River flood plains. The samples were analyzed for physical properties and chemical constituents that included particle-size distribution, water content, volatile solids, nutrients, carbon, selected trace elements, pesticides, and semivolatile organic compounds.

Postflood sediment samples consist primarily of silt and clay; sand ranged from 0 to 28.4 percent by weight in the samples. Moisture content ranged from 7 to 51 percent by weight. Volatile solids concentrations ranged from 6,750 to 91,500 milligrams per kilogram. Total carbon was present predominantly as organic carbon in postflood samples. Several pesticides commonly applied to cropland were detected in samples from several sites. Alachlor was detected in 14 samples; the largest concentration detected was 54 micrograms per kilogram. Atrazine and 4,4'-DDE were detected in 13 samples. Dieldrin was detected in seven samples, all from the Missouri River flood plain. Metolachlor was detected in 15 samples. Eight semivolatile organic compounds were detected in samples from 8 sites in the Missouri River flood plain; 25 semivolatile organic compounds were detected in samples from 7 sites in the Mississippi River flood plain.

REFERENCES CITED

- Arbogast, B.F., ed., 1990, Quality assurance manual for the Branch of Geochemistry, U.S. Geological Survey: U.S. Geological Survey Open-File Report 90-668, 184 p.
- Fishman, M.J., ed., 1993, Method of analysis by the U.S. Geological Survey National Water Quality Laboratory—Determination of inorganic and organic constituents in water and fluvial sediments: U.S. Geological Survey Open-File Report 93-125, 217 p.
- Fishman, M.J., and Friedman, L.C., eds., 1989, Methods for determination of inorganic substances in water and fluvial sediments: U.S. Geological Survey Techniques of Water-Resources Investigations, book 5, chap. A1, 545 p.
- Guy, H.P., 1969, Laboratory techniques and methods for sediment analysis: U.S. Geological Survey Techniques of Water-Resources Investigations, book 5, chap. C1, 58 p.
- Gladney, E.S., Burns, C.E., and Roelandts, I., 1984, 1982 compilation of elemental concentration data for the United States Geological Survey's geochemical explo-

- ration reference samples GXR-1 to GXR-6: Geostandards Newsletter, v. 8, no. 2, p. 119–154.
- Horowitz, A.J., 1991, A primer on sediment-trace element chemistry (2d ed.): Chelsea, Mich., Lewis Publishers, 136 p.
- National Institute of Standards and Technology, 1990, Certificate of analysis, standard reference material 1941: Office of Standard Reference Materials, 11 p.
- O'Leary, R.M., and Viets, J.G., 1986, Determination of antimony, arsenic, bismuth, cadmium, copper, lead, molybdenum, silver, and zinc in geological materials by atomic absorption spectrometry using a hydrochloric acid-hydrogen peroxide digestion: Atomic Spectroscopy, v. 7, p. 4–8.
- Parrett, Charles, Melcher, N.B., and James, R.W., Jr., 1993, Flood discharges in the Upper Mississippi River Basin, 1993: U.S. Geological Survey Circular 1120-A, 14 p.
- U.S. Army Corps of Engineers, 1970, Grain-size analyses, appen. V of Engineer manual—Laboratory soils testing, engineering and design: U.S. Army Corp of Engineers, EM 1110-2-1906, 28 p.
- U.S. Environmental Protection Agency, 1996, Test methods for evaluating solid waste, v. 1B of Laboratory manual physical/chemical methods (3d ed.): Office of Solid Waste and Emergency Response, SW-846, p. 8080–1—8080–27 and 8270–1—8270–34.
- Wershaw, R.L., Fishman, M.J., Grabbe, R.R., and Lowe, L.E., eds., 1987, Methods for the determination of organic substances in water and fluvial sediments: U.S. Geological Survey Techniques of Water-Resources Investigations, book 5, chap. A3, 80 p.
- Xie, Xuejing, and Yan, Mingcai, 1985, Geochemical reference samples, drainage sediment GSE 1–8 from China: Geostandards Newsletter, v. 9, no. 1, p. 83–159.

EXPLANATION OF TABLE 7

ABBREVIATIONS AND REPORTING UNITS FOR CHEMICAL CONSTITUENTS AND NOTATIONS USED IN TABLE

wt. %	Percent by weight
mg/kg	Milligrams per kilogram
4N HCl	4 normal hydrochloric acid (conditionally available fraction)
0.1N HCl	0.1 normal hydrochloric acid (available fraction)
AAS	Atomic absorption spectrometry
CVAAS	Cold-vapor atomic absorption spectrometry
DN	Delayed neutron activation
CO ₂	Carbon dioxide
MO	Missouri
MS	Mississippi
rep	Replicate sample
dup	Duplicate sample
—	No data
N	Not detected at the lower reporting limit
L	Detected at less than the lower reporting limit but not quantified

All elements analyzed by inductively coupled plasma-atomic emission spectrometry except where noted. Analyzed at the U.S. Geological Survey laboratory, Lakewood, Colorado.

Table 7. Concentrations of elements and carbon in the less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains

Site no. (fig. 1)	Aluminum			Calcium			Iron			Magnesium		
	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)
Missouri River												
MO.1	5.1	4,300	360	1.4	1,000	8,000	1.8	8,600	620	0.65	1,900	1,900
MO.2	5.7	5,500	480	1.5	1,400	10,000	2.3	11,000	860	.77	2,300	2,100
MO.3	6.8	12,000	1,000	1.2	1,000	8,600	3.4	19,000	1,600	.84	3,000	1,500
¹ MO.4	4.9	3,900	350	1.2	760	7,000	1.6	7,800	660	.51	1,500	1,400
² MO.4	4.3	—	—	2.0	—	—	2.1	—	—	.79	—	—
MO.5	5.6	5,600	510	1.1	820	7,600	2.4	11,000	710	.57	1,700	1,200
MO.6	7.9	9,100	1,400	1.3	1,000	10,000	4.3	15,000	1,900	1.0	2,500	1,400
MO.7	5.3	5,600	430	1.3	960	8,400	2.0	9,900	830	.62	2,100	1,400
MO.7 rep	5.3	5,700	390	1.3	1,000	7,800	1.9	10,000	740	.59	2,200	1,300
MO.7 rep	5.3	4,900	400	1.3	100	8,100	1.9	9,100	730	.59	1,900	1,400
MO.8	7.4	15,000	1,400	1.1	1,100	8,700	3.9	22,000	1,400	.92	3,400	1,600
MO.9	7.9	19,000	1,300	1.1	1,200	8,700	4.3	29,000	1,700	1.0	4,200	1,700
MO.10	5.1	5,800	400	1.0	820	5,700	1.8	10,000	650	.54	2,000	1,200
MO.11	5.8	7,900	600	1.2	1,000	7,600	2.4	14,000	940	.69	2,700	1,500
MO.12	5.6	7,100	570	1.3	1,200	8,400	2.2	13,000	590	.71	2,700	1,700
Mississippi River												
MS.1	5.1	5,900	580	1.4	810	7,900	2.6	13,000	2,200	.69	1,800	2,400
MS.1 rep	5.1	5,600	610	1.4	790	8,800	2.5	13,000	2,100	.70	1,700	2,600
MS.1 dup	5.1	5,600	560	1.4	850	8,100	2.5	13,000	1,900	.69	1,800	2,400
MS.2	6.6	10,000	980	1.0	740	6,800	3.7	20,000	2,100	.77	2,600	1,700
MS.3	4.7	4,600	450	1.7	1,200	10,000	2.0	10,000	1,400	.70	1,700	2,900
MS.4	5.8	8,500	780	2.0	1,300	15,000	3.1	17,000	1,600	.80	2,300	2,400
MS.5	6.1	9,000	710	1.0	680	7,000	3.0	16,000	1,700	.64	2,000	1,400
MS.6	5.2	6,000	440	.8	440	4,400	2.1	10,000	650	.38	1,200	450
MS.7	6.0	8,800	790	1.5	1,100	12,000	3.0	17,000	2,100	.70	2,200	2,000
MS.7 rep	5.9	8,000	740	1.6	1,000	12,000	3.0	15,000	2,000	.70	2,000	1,900
MS.7 dup	5.9	7,500	730	1.6	920	12,000	3.0	14,000	1,800	.70	1,800	2,000
MS.8	4.8	5,000	430	.9	430	5,300	1.6	7,700	440	.38	960	780
MS.9	5.9	8,100	480	1.3	1,100	8,300	2.4	15,000	700	.73	2,900	1,600
MS.10	4.6	4,300	320	.7	290	2,300	1.5	7,000	430	.34	1,100	340
MS.11	6.3	8,300	720	1.3	1,100	9,300	2.9	15,000	1,400	.77	2,500	1,700
MS.12	6.7	10,000	940	1.3	1,100	9,600	3.7	18,000	1,700	.88	2,800	1,800
MS.13	4.9	3,800	350	1.3	890	7,400	1.7	8,000	610	.57	1,600	1,800
MS.13 rep	4.8	4,100	390	1.3	950	8,100	1.7	8,500	620	.57	1,700	1,900
MS.13 dup	4.9	3,800	340	1.3	970	7,100	1.7	8,000	580	.58	1,600	1,700

Data on Sediments Deposited in the Missouri and the Mississippi River Flood Plains

Table 7. Concentrations of elements and carbon in the less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Site no. (fig. 1)	Total (wt. %)	Phosphorus		Potassium		Sodium		Sulfur, total, combust- tion (wt. %)		Titanium	
		4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)
Missouri River											
MO.1	.07	150	260	1.8	420	7.5N	1.1	42	25	.05L	.25
MO.2	.08	230	250	1.8	620	7.5N	.97	40	32	.05L	.27
MO.3	.10	500	130	1.8	1,400	120	.74	54	51	.05L	.30
MO.4	.06	150	260	1.7	290	7.5N	1.1	31	34	.05L	.24
MO.4 ²	.09	—	—	1.6	—	—	1.2	—	—	.05L	.57
MO.5	.07	230	180	1.7	440	7.5N	.89	30	43	.05L	.26
MO.6	.11	360	95	2.0	1,400	460	.47	35	61	.06	.31
MO.7	.07	200	280	1.9	700	7.5N	1.1	35	26	.05L	.24
MO.7 rep	.07	160	300	1.8	750	7.5N	1.1	38	24	.05L	.24
MO.7 dup	.07	190	300	1.9	580	7.5N	1.1	34	21	.05L	.24
MO.8	.10	450	120	1.9	1,800	290	.61	49	60	.05L	.31
MO.9	.11	490	120	1.9	2,400	300	.52	46	73	.05	.32
MO.10	.06	170	260	1.8	680	7.5N	1.1	38	25	.05L	.24
MO.11	.08	320	260	1.9	1,100	7.5N	.99	41	32	.05L	.25
MO.12	.08	210	280	1.9	1,100	7.5N	1.1	58	54	.05L	.23
Mississippi River											
MS.1	.10	450	180	1.8	94	7.5N	1.0	28	28	.05L	.25
MS.1 rep	.09	430	190	1.8	68	7.5N	1.0	25	35	.05	.25
MS.1 dup	.09	480	200	1.7	64	7.5N	.98	22	33	.05	.24
MS.2	.14	750	140	1.8	670	7.5N	.70	27	33	.06	.30
MS.3	.08	360	190	1.6	43	7.5N	1.0	16	16	.05L	.25
MS.4	.12	620	130	1.7	430	7.5N	.74	22	30	.08	.28
MS.5	.11	550	150	1.6	550	7.5N	.69	17	13	.05L	.28
MS.6	.08	250	240	1.6	390	7.5N	.80	9.5	6.7	.05L	.28
MS.7	.09	480	96	1.7	490	7.5N	.74	20	33	.08	.29
MS.7 rep	.09	410	87	1.7	400	7.5N	.76	20	32	.09	.26
MS.7 dup	.09	430	86	1.7	340	7.5N	.76	21	43	.09	.25
MS.8	.04	170	60	1.8	170	7.5N	.80	13	23	.05L	.25
MS.9	.07	290	210	1.9	1,100	7.5N	.94	49	33	.05L	.27
MS.10	.04	150	77	1.8	430	7.5N	.96	19	20	.05L	.26
MS.11	.09	410	170	1.8	940	7.5N	.81	32	32	.05	.27
MS.12	.11	530	130	1.8	1,100	33	.66	37	44	.05	.32
MS.13	.07	180	290	1.7	220	7.5N	1.1	28	21	.05L	.25
MS.13 rep	.07	180	310	1.7	270	7.5N	1.1	30	21	.05L	.29
MS.13 dup	.07	200	310	1.7	210	7.5N	1.1	27	33	.05L	.27

Table 7. Concentrations of elements and carbon in the less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Site no. (fig. 1)	Antimony			Arsenic			Barium		
	Total, AAS (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total, AAS (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)
Missouri River									
MO.1	0.8	6.3N	6.3N	10L	3.8N	3.8N	7.2	3.6	1.2
MO.2	.8	6.3N	6.3N	10	3.8N	3.8N	10	4.3	1.9
MO.3	.9	6.3N	6.3N	14	3.8N	3.8N	14	8.8	1.5
MO.4	.5	6.3N	6.3N	10L	3.8N	3.8N	6.4	2.9	1.3
2MO.4	.6	—	—	10L	—	—	—	—	—
MO.5	.7	6.3N	6.3N	10L	3.8N	3.8N	9.0	4.6	.9
MO.6	1.1	6.3N	6.3N	18	6.9	3.8N	19	6.9	2.0
MO.7	.9	6.3N	6.3N	10L	3.8N	3.8	8.2	3.3	1.9
MO.7 rep	.6	6.3N	6.3N	10L	3.8N	3.8N	8.0	3.5	1.6
MO.7 dup	.8	6.3N	6.3N	10L	3.8N	3.8N	8.1	3.2	2.1
MO.8	1.4	6.3N	6.3N	17	4.2	3.8N	16	9.4	1.6
MO.9	1.2	6.3N	6.3N	22	3.9	3.8N	18	11	1.8
MO.10	.5	6.3N	6.3N	10L	3.8N	3.8	7.0	3	1.7
MO.11	.8	6.3N	6.3N	10	4.0	3.8N	9.8	5.0	2.0
MO.12	.7	6.3N	6.3N	10L	3.8N	3.8N	9.0	4.7	1.6
Mississippi River									
MS.1	0.4	6.3N	6.3N	10L	3.8N	3.8N	6.4	3.1	1.1
MS.1 rep	.5	6.3N	6.3N	10L	3.8N	3.8N	6.0	2.9	1.0
MS.1 dup	.5	6.3N	6.3N	10L	3.8N	3.8N	6.2	2.4	1.0
MS.2	.7	6.3N	6.3N	11	3.8N	3.8N	11	2.7	1.0
MS.3	1.0	6.3N	6.3N	10L	3.8N	3.8N	4.9	2.1	.7
MS.4	1.0	6.3N	6.3N	10L	3.8N	3.8N	7.6	2.0	.7
MS.5	.9	6.3N	6.3N	10	3.8N	3.8N	9.4	4.3	1.0
MS.6	.6	6.3N	6.3N	10L	3.8N	3.8N	5.0	1.8	.6
MS.7	1.0	6.3N	6.3N	10L	3.8N	3.8N	6.8	2.8	.6
MS.7 rep	.8	6.3N	6.3N	10L	3.8N	3.9	6.8	2.7	.7
MS.7 dup	.9	6.3N	6.3N	10L	3.8N	3.8N	6.5	1.9	.6
MS.8	.7	6.3N	6.3N	10L	3.8N	4.0	4.5	1.6	.4
MS.9	1.0	6.3N	6.3N	11	4.8	4.8	9.5	5.7	1.3
MS.10	.5	6.3N	6.3N	10L	3.8N	3.8N	4.3	1.7	.3
MS.11	1.1	6.3N	6.3N	13	3.8N	3.8N	11	4.0	1.7
MS.12	1.3	6.3N	6.3N	15	3.8N	3.8N	14	6.3	1.4
MS.13	.7	6.3N	6.3N	10L	3.8N	3.8N	5.6	2.2	1.0
MS.13 rep	.8	6.3N	6.3N	10L	3.8N	3.8N	5.9	2.6	1.1
MS.13 dup	.7	6.3N	6.3N	10L	3.8N	3.8N	5.9	1.4	1.0

Data on Sediments Deposited in the Missouri and the Mississippi River Flood Plains

Table 7. Concentrations of elements and carbon in the less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Site no. (fig. 1)	Beryllium			Bismuth			Cadmium			Cerium		
	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)									
Missouri River												
MO.1	1.0	0.056	0.16	10L	7.5N	7.5N	2L	0.5N	0.5N	0.56	59	5.0N
MO.2	1.0	.084	.24	10L	7.5N	7.5N	2L	.5N	.5N	.37	70	5.0N
MO.3	2.0	.17	.45	10L	7.5N	7.5N	2L	.5N	.5N	.48	75	5.0N
MO.4	1.0	.048	.15	10L	7.5N	7.5N	2L	.5N	.5N	.22	59	5.2
MO.4	1.0L	—	—	10L	—	—	2L	—	—	—	140	—
MO.5	1.0	.12	.28	10L	7.5N	7.5N	2L	.5N	.5N	.35	69	5.4
MO.6	2.0	.20	.61	10L	7.5N	7.5N	2L	.5N	.5N	.56	70	5.0N
MO.7	1.0	.084	.21	10L	7.5N	7.5N	2L	.5N	.5N	.32	59	5.0N
MO.7 rep	1.0	.083	.19	10L	7.5N	7.5N	2L	.5N	.5N	.27	57	5.0N
MO.7 dup	1.0	.060	.20	10L	7.5N	7.5N	2L	.5N	.5N	.29	54	5.0N
MO.8	2.0	.25	.53	10L	7.5N	7.5N	2L	.5N	.5N	.51	74	5.0N
MO.9	2.0	.25	.59	10L	7.5N	7.5N	2L	.5N	.5N	.50	79	5.0N
MO.10	1.0	.089	.20	10L	7.5N	7.5N	2L	.5N	.5N	.25	60	5.0N
MO.11	1.0	.084	.27	10L	7.5N	7.5N	2L	.5N	.5N	.37	62	5.0N
MO.12	1.0	.099	.24	10L	7.5N	7.5N	2L	.5N	.5N	.34	55	5.0N
Mississippi River												
MS.1	1.0	0.025N	0.22	10L	7.5N	7.5N	2L	0.5N	0.5N	0.37	54	5.0N
MS.1 rep	1.0	.025N	.24	10L	7.5N	7.5N	2L	.5N	.5N	.40	53	5.0N
MS.1 dup	1.0	.025N	.23	10L	7.5N	7.5N	2L	.5N	.5N	.37	51	5.0N
MS.2	1.0	.064	.43	10L	7.5N	7.5N	2L	.5N	.5N	.40	73	5.0N
MS.3	1.0L	.025N	.17	10L	7.5N	7.5N	2L	.5N	.5N	.35	58	5.0N
MS.4	1.0	.025N	.33	10L	7.5N	7.5N	2L	.5N	.5N	.66	62	5.0N
MS.5	1.0	.063	.39	10L	7.5N	7.5N	2L	.5N	.5N	.27	67	5.0N
MS.6	1.0	.092	.34	10L	7.5N	7.5N	2L	.5N	.5N	.25	67	6.7N
MS.7	1.0	.059	.39	10L	7.5N	7.5N	2L	.5N	.5N	.49	64	5.0N
MS.7 rep	1.0	.053	.36	10L	7.5N	7.5N	2L	.5N	.5N	.52	61	5.0N
MS.7 dup	1.0	.030	.37	10L	7.5N	7.5N	2L	.5N	.5N	.51	65	5.0N
MS.8	1.0	.082	.31	10L	7.5N	7.5N	2L	.5N	.5N	.18	60	5.0N
MS.9	1.0	.11	.26	10L	7.5N	7.5N	2L	.5N	.5N	.35	63	5.0N
MS.10	1.0L	.061	.20	10L	7.5N	7.5N	2L	.5N	.5N	.36	57	5.0N
MS.11	1.0	.11	.37	10L	7.5N	7.5N	2L	.5N	.5N	.50	69	5.0N
MS.12	2.0	.14	.46	10L	7.5N	7.5N	2L	.5N	.5N	.61	71	5.0N
MS.13	1.0	.027	.16	10L	7.5N	7.5N	2L	.5N	.5N	.24	73	5.0N
MS.13 rep	1.0	.026	.16	10L	7.5N	7.5N	2L	.5N	.5N	.26	73	5.0N
MS.13 dup	1.0	.040	.16	10L	7.5N	7.5N	2L	.5N	.5N	.25	76	5.0N

Table 7. Concentrations of elements and carbon in the less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Site no. (fig. 1)	Total (mg/kg)	Chromium		Cobalt		Copper		Europium, total (mg/kg)		Gold, total (mg/kg)		Gali- um, total (mg/kg)		Holmium, total (mg/kg)
		4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	
Missouri River														
MO.1	40	5.2	1.5N	8	1.3	1.7	11	3.1	3.5	2.0L	11	8.0L	4.0L	
MO.2	50	4.8	1.5N	10	2.6	1.3N	17	4.4	5.7	2.0L	12	8.0L	4.0L	
MO.3	65	9.6	1.5N	15	3.4	2.2	25	8.3	8.6	2.0L	16	8.0L	4.0L	
MO.4	35	4.1	1.5N	8	2.0	1.6	10	2.5	2.7	2.0L	10	8.0L	4.0L	
MO.4	55	—	—	—	—	—	7	—	—	2.0L	10	8.0L	4.0L	
MO.5	51	5.1	1.5N	11	2.3	1.8	15	4.8	4.4	2.0L	14	8.0L	4.0L	
MO.6	82	7.1	1.5N	16	2.3	3.2	31	8.2	10	2.0L	21	8.0L	4.0L	
MO.7	41	4.9	1.5N	9	1.7	1.4	14	3.7	4.7	2.0L	12	8.0L	4.0L	
MO.7 rep	42	5.8	1.5N	9	1.8	1.8	14	3.6	4.1	2.0L	11	8.0L	4.0L	
MO.7 dup	40	5.6	1.5N	8	1.5	1.4	14	3.7	4.2	2.0L	11	8.0L	4.0L	
MO.8	73	12	1.5N	16	3.6	2.7	29	11	8.6	2.0L	19	8.0L	4.0L	
MO.9	81	13	1.5N	17	3.9	2.3	31	13	9.4	2.0L	19	8.0L	4.0L	
MO.10	40	6.5	1.5N	9	1.7	1.4	12	4.1	3.6	2.0L	11	8.0L	4.0L	
MO.11	47	6.7	1.5N	11	3.2	1.3	17	6.2	4.9	2.0L	13	8.0L	4.0L	
MO.12	43	6.9	1.5N	10	2.9	2.0	16	6.2	4.2	2.0L	13	8.0L	4.0L	
Mississippi River														
MS.1	50	9.3	1.5N	11	2.0	1.3N	16	3.2	4.8	2.0L	11	8.0L	4.0L	
MS.1 rep	49	8.4	1.5N	11	2.2	1.3N	16	3.2	5.3	2.0L	11	8.0L	4.0L	
MS.1 dup	48	9.3	1.5N	11	1.9	1.3N	16	3.4	4.6	2.0L	11	8.0L	4.0L	
MS.2	67	11	1.5N	15	2.1	3.0	25	7.6	8.4	2.0L	16	8.0L	4.0L	
MS.3	43	6.8	1.5N	8	N1.3	1.3	14	3.9	5.2	2.0L	10	8.0L	4.0L	
MS.4	62	11	1.5N	12	2.2	1.8	21	6.1	6.4	2.0L	14	8.0L	4.0L	
MS.5	60	8.6	1.5N	12	2.5	1.6	22	6.5	6.6	2.0L	13	8.0L	4.0L	
MS.6	47	6.6	1.5N	8	1.7	1.3N	17	4.8	5.1	2.0L	12	8.0L	4.0L	
MS.7	59	9.5	1.5N	12	2.1	2.2	21	5.6	8.1	2.0L	13	8.0L	4.0L	
MS.7 rep	57	9.5	1.7	12	2.5	2.8	20	5.5	7.5	2.0L	13	8.0L	4.0L	
MS.7 dup	59	9.2	1.6	12	2.5	2.8	21	5.2	7.2	2.0L	13	8.0L	4.0L	
MS.8	40	6.3	1.5N	6	1.4	1.3N	15	4.0	4.5	2.0L	11	8.0L	4.0L	
MS.9	52	8.5	1.5N	11	2.9	2.2	18	6.6	5.4	2.0L	14	8.0L	4.0L	
MS.10	38	4.2	1.6	7	2.1	1.7	14	3.3	3.6	2.0L	9	8.0L	4.0L	
MS.11	59	8.2	1.5N	12	3.1	2.5	21	6.5	7.6	2.0L	15	8.0L	4.0L	
MS.12	74	9.6	1.6	15	3.3	3.4	26	7.7	9.2	2.0L	19	8.0L	4.0L	
MS.13	43	4.6	1.5N	7	1.7	1.3	12	2.4	2.6	2.0L	11	8.0L	4.0L	
MS.13 rep	40	5.2	1.5N	8	1.6	1.3N	11	2.5	2.6	2.0L	11	8.0L	4.0L	
MS.13 dup	40	4.8	1.7	8	1.6	1.7	11	2.6	2.8	2.0L	11	8.0L	4.0L	

Data on Sediments Deposited in the Missouri and the Mississippi River Flood Plains

Table 7. Concentrations of elements and carbon in the less than 0.062-millimeter size fraction of postflood sediment samples in the Mississippi and the Missouri River flood plains—Continued

Site no. (fig. 1)	Lanthanum			Lead			Lithium			Manganese		
	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)									
Missouri River												
MO.1	35	7.7	2.6	15	6.3N	21	5.0	0.38N	0.38N	4.30	84	210
MO.2	39	8.8	3.2	17	6.3N	9.2	6.1	.38N	.38N	650	120	380
MO.3	43	13	5.7	22	7.0	8.6	37	9.9	9.9	1,100	370	560
¹ MO.4	35	7.4	2.6	16	6.3N	6.3N	19	4.4	4.4	450	93	240
² MO.4	78	—	—	13	—	—	14	—	—	500	—	—
MO.5	37	9.6	3.6	16	11	6.7	28	5.6	5.6	760	280	320
MO.6	40	10	6.2	24	12	10	46	9.0	9.0	1,300	330	720
MO.7	35	8.4	3.0	18	7.0	6.3N	22	5.6	5.4	490	89	280
MO.7 rep	34	8.9	3.0	17	6.3N	6.4	22	4.9	4.9	460	94	240
MO.7 dup	33	8.1	3.3	16	6.3N	6.3N	22	4.9	4.9	450	87	250
MO.8	44	16	6.1	24	9.8	9.5	41	11	11	1,300	470	640
MO.9	45	18	6.6	23	12	8.1	45	14	14	1,200	490	610
MO.10	36	9.7	3.2	15	7.4	6.9	21	5.0	5.0	450	180	180
MO.11	36	11	3.9	15	7.3	6.8	25	6.7	6.7	820	380	320
MO.12	32	10	3.4	17	12	6.3N	23	6.5	6.5	770	330	320
Mississippi River												
MS.1	30	6.5	4.0	19	9.1	9.4	20	4.1	4.1	0.38N	1,100	400
MS.1 rep	30	6.4	4.1	21	12	8.5	20	3.8	3.8	.38N	1,100	340
MS.1 dup	29	6.7	3.9	21	6.3N	6.3N	20	4.3	4.3	.38N	1,100	360
MS.2	40	11	6.7	22	11	14	31	7.2	7.2	.38N	1,400	280
MS.3	33	5.6	2.8	23	6.7	13	17	3.4	3.4	.38N	660	110
MS.4	35	8.0	4.6	23	15	11	29	6.5	6.5	.38N	1,200	230
MS.5	39	11	5.5	18	12	6.3N	30	6.2	6.2	.38N	970	380
MS.6	38	9.9	4.4	16	7.9	6.3N	24	3.8	3.8	.38N	570	270
MS.7	36	9.4	6.0	24	11	10	29	6.4	6.4	.38N	1,100	170
MS.7 rep	34	8.8	5.7	21	13	9.9	29	6.1	6.1	.38N	1,000	150
MS.7 dup	35	8.3	5.6	22	6.3N	6.3N	29	5.6	5.6	.38N	1,000	140
MS.8	35	8.1	4.8	19	17	6.3N	22	3.4	3.4	.38N	830	420
MS.9	36	12	3.8	18	12	6.3N	28	8.7	8.7	.38N	750	270
MS.10	33	5.8	3.3	21	17	6.3N	17	3.5	3.5	.38N	610	320
MS.11	40	11	5.3	22	6.3N	6.3N	31	7.6	7.6	.38N	970	280
MS.12	39	12	6.4	25	6.3N	6.3N	40	9.0	9.0	.38N	1,300	300
MS.13	41	7.0	2.9	16	6.3N	6.3N	19	3.9	3.9	.38N	640	220
MS.13 rep	41	7.8	3.0	16	6.3N	6.3N	19	4.2	4.2	.38N	670	230
MS.13 dup	43	7.2	3.1	15	6.3N	6.3N	19	4.0	4.0	.38N	670	220

Table 7. Concentrations of elements and carbon in the less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Site no. (fig. 1)	Mercury (CVAAS) Total (mg/kg)	Molybdenum			Neody- mium, total (mg/kg)	Nickel Total (mg/kg)	Niobium, total (mg/kg)	Scan- dium, total (mg/kg)
		0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)			
Missouri River								
MO.1	0.02L	0.008	2.0L	1.0N	1.0N	26	18	4.0
MO.2	.02L	.006	2.0L	1.0N	1.0N	31	24	7.7
MO.3	.02	.010	2.0L	1.5	1.0N	34	33	3.5
¹ MO.4	.02L	.006L	2.0L	1.0N	1.0N	27	16	6.1
² MO.4	.02L	—	2.0L	—	—	60	14	12
MO.5	.02L	.006	2.0L	1.1	1.0N	30	24	6.8
MO.6	.03	.010	2.0L	1.0N	1.0N	34	40	8.3
MO.7	.02L	.006L	2.0L	1.0N	1.0N	30	20	6.4
MO.7 rep	.02L	.006	2.0L	1.0N	1.0N	27	19	7.4
MO.7 dup	.02L	.006L	2.0L	1.1	1.0N	26	19	8.3
MO.8	.02	.006	2.0L	1.0N	1.0N	37	37	4.1
MO.9	.02L	.006	2.0L	1.0N	1.0N	36	42	13
MO.10	.02L	.006L	2.0L	1.0N	1.0N	28	18	6.4
MO.11	.02L	.020L	2.0L	1.0N	1.0N	29	25	3.4
MO.12	.02L	.060L	2.0L	1.0N	1.0N	25	22	11
Mississippi River								
MS.1	0.03	0.008	2.0L	1.0N	1.0N	24	21	5.9
MS.1 rep	.03	.008	2.0L	1.0N	1.0N	21	21	6.5
MS.1 dup	.03	.01	2.0L	1.0N	1.0N	23	22	5.5
MS.2	.04	.006	2.0L	1.0N	1.0N	31	33	1.3N
MS.3	.02L	.006L	2.0L	1.0N	1.0N	28	20	5.6
MS.4	.04	.006	2.0L	1.5	1.0N	28	28	1.3N
MS.5	.02	.006L	2.0L	1.7	1.0N	31	27	8.0
MS.6	.02L	.006L	2.0L	1.4	1.0N	32	21	2.8
MS.7	.02L	.006	2.0L	1.0N	1.0N	29	27	2.1
MS.7 rep	.03	.006L	2.0L	1.4	1.0N	28	27	2.9
MS.7 dup	.04	.01	2.0L	1.7	1.0N	29	28	10
MS.8	.02L	.006L	2.0L	1.1	1.0N	30	18	1.6
MS.9	.02L	.006L	2.0L	1.5	1.0N	30	25	2.0
MS.10	.02L	.006L	2.0L	1.0N	1.0N	26	15	1.1
MS.11	.02	.006L	2.0L	1.1	1.0N	31	28	1.5
MS.12	.03	.006	2.0L	1.0	1.0N	34	36	3.3
MS.13	.02L	.006L	2.0L	1.0N	1.0N	32	17	1.3N
MS.13 rep	.02L	.006L	2.0L	1.1	1.0N	33	17	6.9
MS.13 dup	.02L	.006L	2.0L	1.0N	1.0N	26	16	1.3N

Data on Sediments Deposited in the Missouri and the Mississippi River Flood Plains

Table 7. Concentrations of elements and carbon in the less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Site no. (fig.1)	Selenium (AAS)			Silver			Strontium			Tantala- tum, total (mg/kg)			Thorium		
	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total AAS (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	DN (mg/kg)	
Missouri River															
MO.1	0.3	0.05L	0.05L	2.0L	0.50N	0.50N	0.15	190	9.7	17	40L	8	8.9		
MO.2	.5	.05L	.05L	2.0L	.50N	.50N	.17	180	11	24	40L	10	10		
MO.3	.8	.05L	.05L	2.0L	.50N	.50N	.20	150	14	30	40L	11	14		
MO.4	.2	.05L	.05L	2.0L	.50N	.50N	.14	180	8.1	18	40L	9	8.0		
MO.4	.1	—	—	2.0L	—	—	.11	220	—	—	40L	25	—		
MO.5	.5	.06	.05L	2.0L	.50N	.50N	.16	160	8.7	25	40L	10	9.1		
MO.6	1.0	.05L	.05L	2.0L	.50N	.50N	.17	140	12	46	40L	13	13		
MO.7	.4	.05L	.05L	2.0L	.50N	.50N	.17	180	10	23	40L	11	7.8		
MO.7 rep	.3	.05L	.05L	2.0L	.50N	.50N	.16	180	11	21	40L	9	9.6		
MO.7 dup	.3	.05L	.05L	2.0L	.50N	.50N	.16	180	10	22	40L	9	12		
MO.8	.8	.05L	.05L	2.0L	.50N	.50N	.17	140	15	35	40L	13	12		
MO.9	1.0	.05L	.05L	2.0L	.50N	.50N	.16	140	17	39	40L	13	11		
MO.10	.3	.05L	.05L	2.0L	.50N	.50N	.15	170	9.4	17	40L	9	8.1		
MO.11	.6	.05L	.05L	2.0L	.50N	.50N	.15	170	11	23	40L	9	10		
MO.12	.5	.05L	.05L	2.0L	.50N	.50N	.16	190	13	24	40L	8	7.3		
Mississippi River															
MS.1	0.5	0.05L	0.05L	2.0L	0.50N	0.50N	0.16	150	4.5	10	40L	8	7.2		
MS.1 rep	.5	.05L	.05L	2.0L	.50N	.50N	.16	150	4.3	11	40L	8	7.8		
MS.1 dup	.5	.05L	.05L	2.0L	.50N	.50N	.19	150	4.3	10	40L	8	7.9		
MS.2	.8	.05L	.05L	2.0L	.50N	.50N	.16	120	6.3	13	40L	11	10		
MS.3	.3	.05L	.05L	2.0L	.50N	.50N	.13	150	4.0	40L	9	6.9			
MS.4	.8	.05L	.05L	2.0L	.50N	.50N	.21	120	5.6	15	40L	9	8.7		
MS.5	.6	.05L	.05L	2.0L	.50N	.50N	.16	110	6.2	13	40L	10	11		
MS.6	.4	.05L	.05L	2.0L	.50N	.50N	.15	120	4.2	13	40L	9	12		
MS.7	.7	.05L	.05L	2.0L	.50N	.50N	.22	120	6.3	15	40L	11	10		
MS.7 rep	.7	.05L	.05L	2.0L	.50N	.50N	.20	120	5.9	14	40L	10	11		
MS.7 dup	.7	.05L	.05L	2.0L	.50N	.50N	.20	120	5.7	14	40L	9	11		
MS.8	.3	.05L	.05L	2.0L	.50N	.50N	.15	120	3.4	8.8	40L	10	9.6		
MS.9	.4	.05L	.05L	2.0L	.50N	.50N	.17	170	14	23	40L	9	11		
MS.10	.3	.05L	.05L	2.0L	.50N	.50N	.13	140	4.5	7.5	40L	9	9.2		
MS.11	.6	.05L	.05L	2.0L	.50N	.50N	.20	150	11	23	40L	12	12		
MS.12	.8	.05L	.05L	2.0L	.50N	.50N	.23	140	11	24	40L	12	13		
MS.13	.2	.05L	.05L	2.0L	.50N	.50N	.15	180	7.3	13	40L	10	10		
MS.13 rep	.2	.05L	.05L	2.0L	.50N	.50N	.13	170	7.7	14	40L	11	13		
MS.13 dup	.2	.05L	.05L	2.0L	.50N	.50N	.15	180	7.1	13	40L	10	13		

Table 7. Concentrations of elements and carbon in the less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Site no. (fig. 1)	Total (mg/kg)	Tin		Tungsten		Uranium		Vanadium		Ytter- bium, total (mg/kg)
		4N HCl (mg/kg)	0.1N HCl (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	Total DN (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	
Missouri River										
MO.1	5.0L	10N	10N	7.5N	7.5N	100L	3.1	63	12	3.2
MO.2	5.0L	10N	10N	7.5N	7.5N	100L	3.3	79	14	5.1
MO.3	5.0L	10N	10N	7.5N	7.5N	100L	3.7	110	26	8.4
¹ MO.4	5.0L	10N	10N	7.5N	7.5N	100L	3.2	56	11	3.0
² MO.4	5.0L	—	—	—	—	100L	—	75	—	—
MO.5	5.0L	10N	10N	7.5N	7.5N	100L	3.6	81	17	4.3
MO.6	5.0L	10N	10N	7.5N	7.5N	100L	3.5	140	20	10
MO.7	5.0L	10N	10N	7.5N	7.5N	100L	3.3	68	13	4.6
MO.7 rep	5.0L	10N	10N	7.5N	7.5N	100L	3.0	65	14	4.0
MO.7 dup	5.0L	10N	10N	7.5N	7.5N	100L	2.9	64	12	3.7
MO.8	5.0L	10N	10N	7.5N	7.5N	100L	3.7	130	33	8.6
MO.9	5.0L	10N	10N	7.5N	7.5N	100L	3.6	140	38	11
MO.10	5.0L	10N	10N	7.5N	7.5N	100L	3.3	64	14	4.7
MO.11	5.0L	10N	10N	7.5N	7.5N	100L	3.3	83	19	5.1
MO.12	5.0L	10N	10N	7.5N	7.5N	100L	2.8	75	18	3.7
Mississippi River										
MS.1	5.0L	10N	10N	7.5N	7.5N	100L	2.6	63	13	4.3
MS.1 rep	5.0L	10N	10N	7.5N	7.5N	100L	2.5	63	13	4.5
MS.1 dup	5.0L	10N	10N	7.5N	7.5N	100L	2.6	62	13	3.7
MS.2	5.0L	10N	10N	7.5N	7.5N	100L	3.2	100	23	9.5
MS.3	5.0L	10N	10N	7.5N	7.5N	100L	2.9	54	11	3.6
MS.4	5.0L	10N	10N	7.5N	7.5N	100L	2.8	83	17	6.4
MS.5	5.0L	10N	10N	7.5N	7.5N	100L	3.3	88	20	6.7
MS.6	5.0L	10N	10N	7.5N	7.5N	100L	3.1	65	12	6.5
MS.7	5.0L	10N	10N	7.5N	7.5N	100L	3.1	86	17	8.6
MS.7 rep	5.0L	10N	10N	7.5N	7.5N	100L	2.9	83	16	8.1
MS.7 dup	5.0L	10N	10N	7.5N	7.5N	100L	3.0	83	15	7.5
MS.8	5.0L	10N	10N	7.5N	7.5N	100L	2.9	57	12	3.9
MS.9	5.0L	10N	10N	7.5N	7.5N	100L	3.0	84	21	4.5
MS.10	5.0L	10N	10N	7.5N	7.5N	100L	3.3	56	12	4.4
MS.11	5.0L	10N	10N	7.5N	7.5N	100L	3.3	93	18	7.3
MS.12	5.0L	10N	10N	7.5N	7.5N	100L	3.4	110	22	9.2
MS.13	5.0L	10N	10N	7.5N	7.5N	100L	3.7	57	11	2.2
MS.13 rep	5.0L	10N	10N	7.5N	7.5N	100L	4.0	60	12	2.1
MS.13 dup	5.0L	10N	10N	7.5N	7.5N	100L	3.9	59	11	2.2

Data on Sediments Deposited in the Missouri and the Mississippi River Flood Plains

Table 7. Concentrations of elements and carbon in the less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Site no. (fig. 1)	Yttrium			Zinc			Zirconium			Carbon (wt. %)		
	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total, combustion	Organic, diff.- ference	Car- bonate, titration	$^{3}\text{CO}_2$
Missouri River												
MO.1	19	4.2	3.6	51	32	6.4	0.63N	0.63N	0.80	0.051	0.29	1.1
MO.2	21	5.8	4.6	66	40	9.1	.63N	.63N	1.24	.91	.33	1.2
MO.3	24	8.5	6.7	96	64	15	.63N	.63N	1.53	1.37	.16	.57
¹ MO.4	17	3.8	3.3	44	28	6.5	.63N	.63N	.56	.35	.21	.77
² MO.4	28	—	—	42	—	—	—	—	.61	.12	.49	.18
MO.5	19	5.7	4.3	64	35	10	.63N	.63N	.98	.81	.17	.63
MO.6	25	7.6	7.5	130	53	18	.63N	.63N	2.38	2.27	.11	.39
MO.7	18	4.8	4.0	63	37	10	.63N	.63N	.92	.68	.24	.88
MO.7 rep	18	5.1	3.6	57	37	9.7	.63N	.63N	.84	.60	.24	.87
MO.7 dup	17	4.8	3.7	56	37	8.5	.63N	.63N	.90	.67	.23	.86
MO.8	26	1.0	7.4	110	73	14	.63N	.63N	1.85	1.73	.12	.43
MO.9	27	13	7.9	120	86	13	.63N	.63N	1.90	1.81	.09	.35
MO.10	19	5.5	3.7	50	36	5.7	.63N	.63N	.78	.64	.14	.53
MO.11	19	6.9	4.7	70	54	8.8	.63N	.63N	1.19	1.02	.17	.62
MO.12	18	6.5	4.3	64	46	8	.63N	.63N	1.19	.97	.22	.81
Mississippi River												
MS.1	16	3.7	4.3	89	52	22	0.63N	0.63N	1.92	1.65	0.27	0.97
MS.1 rep	16	3.6	4.6	93	52	25	.63N	.63N	2.00	1.73	.27	.98
MS.1 dup	16	3.8	4.2	91	53	24	.63N	.63N	1.95	1.68	.27	.99
MS.2	22	7.0	6.4	100	62	18	.63N	.76	3.10	2.98	.12	.45
MS.3	16	3.0	3.5	67	38	17	.63N	.66	1.54	1.09	.45	1.6
MS.4	20	5.2	5.2	120	66	30	.63N	.63N	3.30	2.86	.44	1.6
MS.5	21	6.3	5.6	83	51	12	.63N	.63N	2.13	2.01	.12	.42
MS.6	21	5.9	4.3	65	39	7.1	.90	.76	1.29	1.26	.03	.10
MS.7	21	5.8	6.2	100	59	24	.63N	.70	2.39	2.08	.31	1.1
MS.7 rep	20	5.4	5.9	98	55	23	.63N	1.1	2.37	2.06	.31	1.1
MS.7 dup	20	5.1	5.7	99	53	25	.63N	.63N	2.54	2.23	.31	1.1
MS.8	18	4.5	4.3	50	29	4.7	.63N	1.8	1.34	1.23	.11	.42
MS.9	20	7.2	4.3	72	54	8.7	.63N	1.4	1.07	.84	.23	.86
MS.10	15	3.0	2.5	44	32	4.4	.63N	2.4	1.19	1.18	.01	.05
MS.11	21	7.0	5.8	88	52	16	.63N	.63N	1.52	1.31	.21	.79
MS.12	23	8.0	6.8	110	62	19	.63N	.80	1.96	1.76	.20	.72
MS.13	17	3.7	3.5	48	28	7.1	.63N	.63N	.70	.46	.24	.89
MS.13 rep	19	3.9	3.7	49	31	8.6	.63N	.64	.38	.26	.94	.94
MS.13 dup	19	3.9	3.3	49	29	7.7	.63N	4.5	.70	.44	.26	.94

¹Layer 1, 0 to 3 centimeters.

²Layer 2, 3 to 15 centimeters.

³Coulometrically titrated to compute carbonate carbon.

EXPLANATION OF TABLE 8

ABBREVIATIONS AND REPORTING UNITS FOR CHEMICAL CONSTITUENTS AND NOTATIONS USED IN TABLE

wt. %	Percent by weight
mg/kg	Milligrams per kilogram
4N HCl	4 normal hydrochloric acid (conditionally available fraction)
0.1N HCl	0.1 normal hydrochloric acid (available fraction)
AAS	Atomic absorption spectrometry
CVAAS	Cold-vapor atomic absorption spectrometry
DN	Delayed neutron activation
MO	Missouri
MS	Mississippi
N	Not detected at the lower reporting limit
L	Detected at less than the lower reporting limit but not quantified
n.a.	Not applicable

All elements analyzed by inductively coupled plasma-atomic emission spectrometry except where noted. Analyzed at the U.S. Geological Survey laboratory, Lakewood, Colorado.

Table 8. Descriptive statistics of selected element and carbon concentrations in less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains

Descriptive statistics	Aluminum			Calcium			Iron			Magnesium		
	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)
Missouri River flood-plain samples (12)												
Minimum.....	4.9	3,900	350	1.0	760	5,700	1.6	7,800	590	0.51	1,500	1,200
Maximum.....	7.9	19,000	1,400	1.5	1,400	10,000	4.3	29,000	1,900	1.0	4,200	2,100
Mean	6.1	8,400	730	1.2	1,000	8,200	2.7	14,000	1,000	.74	2,500	1,600
Standard deviation.....	1.1	4,700	420	.14	180	1,200	1.0	6,300	470	.17	770	270
Number of samples above detection limit	12	12	12	12	12	12	12	12	12	12	12	12
Mississippi River flood-plain samples (13)												
Minimum.....	4.6	3,800	320	0.7	290	2,300	1.5	7,000	430	0.34	960	340
Maximum.....	6.7	10,000	980	2.0	1,300	15,000	3.7	20,000	2,200	.88	2,900	2,900
Mean	5.6	7,100	610	1.2	860	8,100	2.6	13,000	1,300	.64	2,000	1,600
Standard deviation.....	.74	2,200	220	.37	330	3,300	.75	4,400	660	.17	640	750
Number of samples above detection limit	13	13	13	13	13	13	13	13	13	13	13	13
Missouri and Mississippi River flood-plain samples (25)												
Minimum.....	4.6	3,800	320	0.7	290	2,300	1.5	7,000	430	0.34	960	340
Maximum.....	7.9	19,000	1,400	2.0	1,400	15,000	4.3	29,000	2,200	1.0	4,200	2,900
Mean	5.8	7,800	670	1.2	940	8,200	2.6	14,000	1,200	.69	2,200	1,600
Standard deviation.....	.95	3,600	330	.28	270	2,400	.86	5,300	580	.18	740	560
Number of samples above detection limit	25	25	25	25	25	25	25	25	25	25	25	25

Table 8. Descriptive statistics of selected element and carbon concentrations in less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Descriptive statistics	Phosphorus			Potassium			Sodium			Sulfur, total, combustion (wt. %)			Titanium		
	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (wt. %)	4N HCl (mg/kg)	0.1N HCl (mg/kg)
Missouri River flood-plain samples (12)															
Minimum.....	0.06	150	95	1.7	290	7.5N	0.47	30	25	0.05L	0.23	61	0.71		
Maximum.....	.11	500	280	2.0	2,400	460	1.1	58	73	.06	.32	170	2.5		
Mean.....	.08	290	210	1.8	1,000	n.a.	.89	42	43	n.a.	.27	100	1.4		
Standard deviation018	130	73	.09	630	n.a.	.24	8.8	16	n.a.	.032	34	.62		
Number of samples above detection limit.....	12	12	12	12	12	4	12	12	12	2	12	12	12		
Mississippi River flood-plain samples (13)															
Minimum.....	0.04	150	60	1.6	43	7.5N	0.66	9.5	6.7	0.05L	0.25	57	1.2		
Maximum.....	.14	750	290	1.9	1,100	33	1.1	49	44	.08	.32	190	6.5		
Mean.....	.087	400	160	1.7	510	n.a.	.84	24	26	n.a.	.27	120	3.1		
Standard deviation029	190	65	.10	360	n.a.	.14	11	10	n.a.	.022	41	1.7		
Number of samples above detection limit.....	13	13	13	13	1	13	13	13	13	6	13	13	13		
Missouri and Mississippi River flood-plain samples (25)															
Minimum.....	0.04	150	60	1.6	43	7.5N	0.47	9.5	6.7	0.05L	0.23	57	0.71		
Maximum.....	.14	750	290	2.0	2,400	460	1.1	58	73	.08	.32	190	6.5		
Mean.....	.085	340	180	1.8	790	n.a.	.86	33	34	n.a.	.27	110	2.3		
Standard deviation024	170	72	.10	560	n.a.	.19	13	16	n.a.	.027	38	1.5		
Number of samples above detection limit.....	25	25	25	25	5	25	25	25	25	8	25	25	25		

Table 8. Descriptive statistics of selected element and carbon concentrations in less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Descriptive statistics	Antimony				Arsenic				Barium			
	Total, AAS (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total, AAS (mg/kg)	4N HCl, AAS (mg/kg)	0.1N HCl, AAS (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)
Missouri River flood-plain samples (12)												
Minimum	0.5	6.3N	6.3N	10L	3.8N	3.8N	6.4	2.9	0.9	670	53	62
Maximum.....	1.4	6.3N	6.3N	22	6.9	3.8	19	11	2.0	790	120	190
Mean9	n.a.	n.a.	n.a.	n.a.	n.a.	11	5.6	1.6	740	82	110
Standard deviation27	n.a.	n.a.	n.a.	n.a.	n.a.	4.4	2.7	.34	35	20	49
Number of samples above detection limit.....	12	0	0	6	4	2	12	12	12	12	12	12
Mississippi River flood-plain samples (13)												
Minimum	0.4	6.3N	6.3N	10L	3.8N	3.8N	4.3	1.6	0.3	550	24	51
Maximum.....	1.3	6.3N	6.3N	15	4.8	4.8	14	6.3	1.7	770	91	140
Mean84	n.a.	n.a.	n.a.	n.a.	n.a.	7.7	3.1	.91	650	55	97
Standard deviation26	n.a.	n.a.	n.a.	n.a.	n.a.	3.0	1.5	.41	68	23	30
Number of samples above detection limit.....	13	0	0	5	1	2	13	13	13	13	13	13
Missouri and Mississippi River flood-plain samples (25)												
Minimum	0.4	6.3N	6.3N	10L	3.8N	3.8N	4.3	1.6	0.3	550	24	51
Maximum.....	1.4	6.3N	6.3N	22	6.9	4.8	19	11	2.0	790	120	190
Mean8	n.a.	n.a.	n.a..	n.a.	n.a.	9.3	4.4	1.2	690	69	100
Standard deviation26	n.a.	n.a.	n.a.	n.a.	n.a.	4.1	2.5	.52	74	25	40
Number of samples above detection limit.....	25	0	0	11	5	4	25	25	25	25	25	25

Table 8. Descriptive statistics of selected element and carbon concentrations in less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Descriptive statistics	Beryllium			Bismuth			Cadmium			Cerium		
	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)
Missouri River flood-plain samples (12)												
Minimum.....	1	0.048	0.15	10L	7.5N	7.5N	0.5N	0.5N	0.22	55	5.0N	5.0N
Maximum.....	2	.25	.61	10L	7.5N	7.5N	.5N	.5N	.56	79	5.4	10
Mean.....	1.3	.13	.33	n.a.	n.a.	n.a.	n.a.	n.a.	.40	66	n.a.	n.a.
Standard deviation.....	.5	.072	.17	n.a.	n.a.	n.a.	n.a.	n.a.	.12	7.8	n.a.	n.a.
Number of samples above detection limit	12	12	12	0	0	0	0	0	12	12	2	4
Mississippi River flood-plain samples (13)												
Minimum.....	1.0L	0.025N	0.16	10L	7.5N	7.5N	0.5N	0.5N	0.18	54	5.0N	5.0N
Maximum.....	2	.14	.46	10L	7.5N	7.5N	.5N	.5N	.66	73	6.7	12
Mean.....	1	.065	.31	n.a.	n.a.	n.a.	n.a.	n.a.	.39	64	n.a.	n.a.
Standard deviation.....	.4	.041	.10	n.a.	n.a.	n.a.	n.a.	n.a.	.14	6.2	n.a.	n.a.
Number of samples above detection limit	11	10	13	0	0	0	0	1	13	13	2	6
Missouri and Mississippi River flood-plain samples (25)												
Minimum.....	1.0L	0.025N	0.15	10L	7.5N	7.5N	0.5N	0.5N	0.18	54	5.0N	5.0N
Maximum.....	2	.25	.61	10L	7.5N	7.5N	.5N	.5N	.66	79	6.7	12
Mean.....	1.2	.099	.32	n.a.	n.a.	n.a.	n.a.	n.a.	.39	65	n.a.	n.a.
Standard deviation.....	.45	.065	.13	n.a.	n.a.	n.a.	n.a.	n.a.	.13	6.9	n.a.	n.a.
Number of samples above detection limit	23	22	25	0	0	0	0	1	25	25	4	10

Table 8. Descriptive statistics of selected element and carbon concentrations in less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Descriptive statistics	Chromium			Cobalt			Copper			Europium, total (mg/kg)			Gallium, total (mg/kg)			Gold, total (mg/kg)			Holmium, total (mg/kg)		
	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Europlum, total (mg/kg)	Gallium, total (mg/kg)	Gold, total (mg/kg)	Holmium, total (mg/kg)								
Missouri River flood-plain samples (12)																					
Minimum.....	35	4.1	1.5N	8	1.3	1.3N	10	2.5	2.7	2.0L	10	8.0L	4.0L								
Maximum.....	82	13	1.5N	17	3.9	3.2	31	13	10	2.0L	21	8.0L	4.0L								
Mean.....	54	7.2	n.a.	12	2.6	1.8	19	6.3	5.8	n.a.	14	n.a.	n.a.								
Standard deviation.....	17	2.9	n.a.	3.4	.83	.68	7.8	3.2	2.6	n.a.	3.6	n.a.	n.a.								
Number of samples above detection limit.....	12	12	0	12	12	11	12	12	12	0	12	0	0								
Mississippi River flood-plain samples (13)																					
Minimum.....	38	4.2	1.5N	6	1.3N	1.3N	12	2.4	2.6	2.0L	9	8.0L	4.0L								
Maximum.....	74	11	1.6	15	3.3	3.4	26	7.7	9.2	2.0L	19	8.0L	4.0L								
Mean.....	53	8.0	n.a.	10	2.1	1.8	19	5.2	6.0	n.a.	13	n.a.	n.a.								
Standard deviation.....	11	2.2	n.a.	3.0	.72	.88	4.4	1.8	2.0	n.a.	2.7	n.a.	n.a.								
Number of samples above detection limit.....	13	13	2	13	12	10	13	13	13	0	13	0	0								
Missouri and Mississippi River flood-plain samples (25)																					
Minimum.....	35	4.1	1.5N	6	1.3N	1.3N	10	2.4	2.6	2.0L	9	8.0L	4.0L								
Maximum.....	82	13	1.6	17	3.9	3.4	31	13	10	2.0L	21	8.0L	4.0L								
Mean.....	54	7.5	n.a.	11	2.4	1.8	19	5.8	5.9	n.a.	14	n.a.	n.a.								
Standard deviation.....	14	2.5	n.a.	3.2	.79	.78	6.2	2.6	2.2	n.a.	3.2	n.a.	n.a.								
Number of samples above detection limit.....	25	25	2	25	24	21	25	25	25	0	25	0	0								

Table 8. Descriptive statistics of selected element and carbon concentrations in less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Descriptive statistics	Lanthanum			Lead			Lithium			Manganese		
	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)
Missouri River flood-plain samples (12)												
Minimum.....	32	7.4	2.6	15	6.3N	19	4.4	0.38N	430	84	180	
Maximum.....	45	18	6.6	24	12	10	46	14	.38N	1,300	490	720
Mean.....	38	11	4.2	18	7.9	6.6	30	7.4	n.a.	810	270	400
Standard deviation.....	4.1	3.3	1.5	3.6	3.5	2.7	9.9	2.9	n.a.	340	150	180
Number of samples above detection limit.....	12	12	12	12	9	8	12	12	0	12	12	12
Mississippi River flood-plain samples (13)												
Minimum.....	30	5.6	2.8	16	6.3N	6.3N	17	3.4	0.38N	570	110	140
Maximum.....	41	12	6.7	25	17	14	40	9	.38N	1,400	420	870
Mean.....	36	9.0	4.6	20	9.9	n.a.	26	5.7	n.a.	930	280	460
Standard deviation.....	3.3	2.3	1.3	3.0	4.9	n.a.	6.8	2.1	n.a.	280	88	250
Number of samples above detection limit.....	13	13	13	13	10	5	13	13	0	13	13	13
Missouri and Mississippi River flood-plain samples (25)												
Minimum.....	30	5.6	2.6	15	6.3N	6.3N	17	3.4	0.38N	430	84	140
Maximum.....	45	18	6.7	25	17	14	46	14	.38N	1,400	490	870
Mean.....	37	10	4.4	20	8.9	n.a.	28	6.6	n.a.	870	270	430
Standard deviation.....	3.7	2.9	1.4	3.4	4.3	n.a.	8.5	2.6	n.a.	310	120	220
Number of samples above detection limit.....	25	25	25	25	19	13	25	25	0	25	25	25

Table 8. Descriptive statistics of selected element and carbon concentrations in less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Descriptive statistics	Mercury (CVAAS)		Molybdenum		Neodymium		Nickel		Scandium, total (mg/kg)
	Total (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	
Missouri River flood-plain samples (12)									
Minimum.....	0.02L	0.006L	2.0L	1.0N	1.0N	25	16	5.4	3.0
Maximum.....	.03	.01	2.0L	1.5	1.0N	37	42	16	7.3
Mean	n.a.	.008	n.a.	n.a.	n.a.	30	26	9.4	4.7
Standard deviation.....	n.a.	.007	n.a.	n.a.	n.a.	3.9	9.1	3.3	1.5
Number of samples above detection limit.....	3	7	0	2	0	12	12	12	12
Mississippi River flood-plain samples (13)									
Minimum.....	0.02L	0.006L	2.0L	1.0N	1.0N	24	15	4.7	1.3N
Maximum.....	.04	.008	2.0L	1.7	1.0N	34	36	12	5.6
Mean	n.a.	n.a.	n.a.	.95	n.a.	30	24	8.1	2.6
Standard deviation.....	n.a.	n.a.	n.a.	.47	n.a.	2.7	6.3	2.5	1.5
Number of samples above detection limit.....	6	5	0	7	0	13	13	11	13
Missouri and Mississippi River flood-plain samples (25)									
Minimum.....	0.02L	0.006L	2.0L	1.0N	1.0N	24	15	4.7	1.3N
Maximum.....	.04	.01	2.0L	1.7	1.0N	37	42	16	7.3
Mean	n.a.	n.a.	n.a.	n.a.	n.a.	30	25	8.8	3.6
Standard deviation.....	n.a.	n.a.	n.a.	n.a.	n.a.	3.3	7.7	2.9	1.8
Number of samples above detection limit.....	9	12	0	9	0	25	25	23	25

Table 8. Descriptive statistics of selected element and carbon concentrations in less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Descriptive statistics	Selenium (AAS)			Silver			Strontium			Thorium		
	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	Tantalum, total (mg/kg)	Total, DN (mg/kg)
Missouri River flood plain samples (12)												
Minimum.....	0.2	0.05L	0.05L	2.0L	0.5N	0.5N	0.14	140	8.1	17	40L	8
Maximum.....	1.0	.06	.05L	2.0L	.5N	.5N	.20	190	17	46	40L	13
Mean.....	.6	n.a.	n.a.	n.a.	n.a.	n.a.	.16	160	12	27	n.a.	10
Standard deviation.....	.3	n.a.	n.a.	n.a.	n.a.	n.a.	.015	19	2.7	9.1	n.a.	1.9
Number of samples above detection limit.....	12	1	0	0	0	0	12	12	12	0	12	12
Mississippi River flood-plain samples (13)												
Minimum.....	0.2	0.05L	0.05L	2.0L	0.5N	0.5N	0.13	110	3.4	7.5	40L	8
Maximum.....	.8	.05L	.05L	2.0L	.5N	.5N	.23	180	14	24	40L	12
Mean.....	.5	n.a.	n.a.	n.a.	n.a.	n.a.	.17	140	6.8	14	n.a.	9.9
Standard deviation.....	.2	n.a.	n.a.	n.a.	n.a.	n.a.	.033	22	3.2	5.8	n.a.	1.2
Number of samples above detection limit.....	13	0	0	0	0	0	13	13	13	0	13	13
Missouri and Mississippi River flood-plain samples (25)												
Minimum.....	0.2	0.05L	0.05L	2.0L	0.5N	0.5N	0.13	110	3.4	7.5	40L	8
Maximum.....	1.0	.06	.05L	2.0L	.5N	.5N	.23	190	17	46	40L	13
Mean.....	.5	n.a.	n.a.	n.a.	n.a.	n.a.	.17	150	9.3	20	n.a.	10
Standard deviation.....	.2	n.a.	n.a.	n.a.	n.a.	n.a.	.026	25	3.8	9.8	n.a.	1.6
Number of samples above detection limit.....	25	1	0	0	0	0	25	25	25	0	25	25

42 Data on Sediments Deposited in the Missouri and the Mississippi River Flood Plains

Table 8. Descriptive statistics of selected element and carbon concentrations in less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Descriptive statistics	Tin		Tungsten		Uranium		Vanadium		Ytterbium, total (mg/kg)
	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	
Missouri River flood-plain samples (12)									
Minimum.....	5.0L	10N	10N	7.5N	7.5N	100L	2.8	.56	11
Maximum.....	5.0L	10N	10N	7.5N	7.5N	100L	3.7	.140	.38
Mean	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	3.4	.91	20
Standard deviation.....	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	.27	.31	8.6
Number of samples above detection limit.....	0	0	0	0	0	0	12	.12	12
Mississippi River flood-plain samples (13)									
Minimum.....	5.0L	10N	10N	7.5N	7.5N	100L	2.6	.54	11
Maximum.....	5.0L	10N	10N	7.5N	7.5N	100L	3.7	.110	.23
Mean	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	3.1	.77	16
Standard deviation.....	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	.29	.19	4.5
Number of samples above detection limit.....	0	0	0	0	0	0	13	.13	13
Missouri and Mississippi River flood-plain samples (25)									
Minimum.....	5.0L	10N	10N	7.5N	7.5N	100L	2.6	.54	11
Maximum.....	5.0L	10N	10N	7.5N	7.5N	100L	3.7	.140	.38
Mean	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	3.2	.83	18
Standard deviation.....	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	.30	.26	6.8
Number of samples above detection limit.....	0	0	0	0	0	0	25	.25	25

Table 8. Descriptive statistics of selected element and carbon concentrations in less than 0.062-millimeter size fraction of postflood sediment samples in the Missouri and the Mississippi River flood plains—Continued

Descriptive statistics	Yttrium			Zinc			Zirconium			Carbon (wt. %)		
	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	4N HCl (mg/kg)	0.1N HCl (mg/kg)	Total combustion	Organic, difference	Carbo- nate, titration	$^{13}\text{CO}_2$
Missouri River flood-plain samples (12)												
Minimum.....	17	3.8	3.3	44	28	5.7	0.63N	0.63N	0.56	0.35	0.09	0.35
Maximum.....	27	13	7.9	130	86	18	.63N	.63N	2.38	2.27	.33	1.2
Mean.....	21	6.9	5.2	77	49	10	n.a.	n.a.	1.28	1.09	.19	.69
Standard deviation.....	3.5	2.6	1.7	29	18	3.8	n.a.	n.a.	.54	.59	.073	.27
Number of samples above detection limit.....	12	12	12	12	12	0	0	0	12	12	12	12
Mississippi River flood-plain samples (13)												
Minimum.....	15	3.0	2.5	44	28	4.4	0.63N	0.63N	0.70	0.46	0.01	0.05
Maximum.....	23	8.0	6.8	120	66	30	.9	2.4	3.30	3.00	.45	1.64
Mean.....	19	5.4	4.8	80	48	15	n.a.	.84	1.8	1.59	.21	.77
Standard deviation.....	2.6	1.7	1.3	24	13	8.0	n.a.	.65	.77	.74	.14	.50
Number of samples above detection limit.....	13	13	13	13	13	1	8	13	13	13	13	13
Missouri and Mississippi River flood-plain samples (25)												
Minimum.....	15	3.0	2.5	44	28	4.4	0.63N	0.63N	0.56	0.35	0.01	0.05
Maximum.....	27	13	7.9	130	86	30	.9	2.4	3.3	2.98	.45	1.64
Mean.....	20	6.2	5.0	78	48	12	n.a.	n.a.	1.6	1.35	.20	.73
Standard deviation.....	3.1	2.3	1.5	26	15	6.6	n.a.	n.a.	.71	.70	.11	.40
Number of samples above detection limit.....	25	25	25	25	25	1	8	25	25	25	25	25

¹Coulometrically titrated to compute carbonate carbon.

EXPLANATION OF TABLE 11

ABBREVIATIONS AND REPORTING UNITS FOR CHEMICAL CONSTITUENTS AND NOTATIONS USED IN TABLE

MO	Missouri
SRM	Standard reference material
dup	Duplicate sample split in laboratory
rep	Replicate sample
RS	Reagent spiked sample, in percent recovery
MC	Measured concentration
EC	Expected concentration
STD	Standard deviation
<	Less than
—	No data

Detections shown in bold. Analyzed at the U.S. Geological Survey laboratory, Arvada, Colorado.

Table 11. Concentrations of semivolatile organic compounds in the less than 2-millimeter size fraction of postflood sediment samples in the Missouri River flood plain, sample set reagent spike, standard reference material, and surrogate samples

Compound	Site number (fig. 1)											SRM 1941								
	MO.4			Layer 2			MO.7			Sam- ple			Rep	MO.8	MO.10	MO.11	RS	MC	1EC	1STD
	MO.1	MO.2	Layer 1	Sam- ple	Dup	MO.5	Rep	MO.8	MO.10	MO.11	RS	MC								
Concentration, in micrograms per kilogram																				
Acenaphthene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	62	<10	52	2						
Acenaphthylene.....	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	58	260	115	10						
Acridine.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	69	<10	—	—						
Anthracene.....	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	70	370	202	42						
Anthraquinone.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	80	390	—	—						
Azo-benzene.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	66	<20	—	—						
Benz[a]anthracene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	70	580	2550	79						
Benz[b]fluoranthene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	68	720	2780	190						
Benz[k]fluoranthene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	74	520	2444	49						
Benz[g,h,i]perylene.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	66	360	2516	83						
Benz[a]pyrene.....	110	120	100	<10	<10	<10	<10	<10	<10	100	<10	68	780	2670	130					
Benz[c]quinoline.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	75	<10	—	—						
2,2'-Biquinoline	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	54	<20	—	—						
4-Bromophenyl phenyl ether ..	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	68	<10	—	—						
Butylbenzylphthalate	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	63	<10	—	—						
C ₈ -Alkyl-phenol.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	10	56	<10	—	—	—	—	—		
9H-Carbazole	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	10	75	<10	—	—	—	—	—		
bis(2-Chlorothoxy)methane	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	10	52	<10	—	—	—	—	—		
bis(2-Chlorothio)ether	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	10	55	<10	—	—	—	—	—		
bis(2-Chloroisopropyl)ether.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	51	<20	51	<20	—	—	—	—		
4-Chloro-3-methylphenol	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	30	<30	47	<30	—	—	—	—		
2-Chloronaphthalene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	10	60	<10	—	—	—	—	—		
2-Chlorophenol	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	10	55	<10	—	—	—	—	—		
4-Chlorophenyl phenyl ether ..	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	10	67	<10	—	—	—	—	—		
Chrysene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	10	72	660	449	—	—	—	—		
p-Cresol.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	10	10	52	<10	—	—	—	—		
Dibenzo[a,h]anthracene	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	20	107	<20	—	—	—	—	—		
Dibenzothiophene	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	5	69	69	—	—	—	—	—		
Di-n-butylphthalate	54	53	59	<10	<10	<10	<10	<10	<10	<10	60	60	74	380	—	—	—	—		
1,2-Dichlorobenzene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	10	<10	61	<10	—	—	—	—		

Table 11. Concentrations of semivolatile organic compounds in the less than 2-millimeter size fraction of postflood sediment samples in the Missouri River flood plain, sample set reagent spike, standard reference material, and surrogate samples—Continued

Compound	Site number (fig. 1)										¹ STD	
	MO.4					MO.7						
	Layer 2		Sam- ple	Dup	MO.5	Rep	MO.8	MO.10	MO.11	RS		
Concentration, in micrograms per kilogram—Continued												
1,3-Dichlorobenzene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	58	<10	
1,4-Dichlorobenzene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	49	<10	
2,4-Dichlorophenol.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	56	<20	
Diethyl phthalate.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	70	<10	
1,2-Dimethyl naphthalene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	61	110	
1,6-Dimethyl naphthalene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	61	280	
2,6-Dimethyl naphthalene.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	60	240	
3,5-Dimethylphenol.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	66	<10	
Dimethylphthalate.....	<200	<200	<200	<200	<200	<200	<200	<200	<200	0	<200	
4,6-Dinitro-2-methylphenol.....	<300	<300	<300	<300	<300	<300	<300	<300	<300	0	<300	
2,4-Dinitrophenol.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	59	<20	
2,4-Dinitrotoluene.....	<50	<50	<50	<50	<50	<50	<50	<50	<50	58	<50	
2,6-Dinitrotoluene.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	65	<20	
di-n-Octyl phthalate.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	79	1,900	
bis(2-Ethylhexyl)phthalate.....	60	79	94	<10	96	91	81	66	<10	64	<10	
2-Ethylaphthalene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	58	110	
Fluoranthene.....	36	47	<5	44	58	38	<5	42	77	950	2,1220	
9H-Fluorene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	64	240	
Hexachlorobenzene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	68	5	
Hexachlorobutadiene.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	61	<20	
Hexachlorocyclopentadiene.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	34	<20	
Hexachloroethane.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	51	<20	
Indeno[1,2,3-cd]pyrene.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	52	2569	
Isophorone.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	76	40	
Isoquinoline.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	54	<10	
1-Methyl-9H-fluorene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	68	<10	
2-Methylnaphthalene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	72	320	
4,5-Methyleneephantrene.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	73	260	
1-Methylphenanthrene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	73	109	
1-Methylpyrene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	74	340	

Table 11. Concentrations of semivolatile organic compounds in the less than 2-millimeter size fraction of postflood sediment samples in the Missouri River flood plain, sample set reagent spike, standard reference material, and surrogate samples—Continued

Compound	Site number (fig. 1)										SRM 1941		
	MO.4			MO.7			MO.11						
	MO.1	MO.2	Layer 1	Sam- ple	Dup	MO.5	Rep	MO.8	MO.10	MO.11			
Concentration, in micrograms per kilogram—Continued													
Naphthalene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	58	780	
Nitrobenzene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	52	<10	
2-Nitrophenol	<100	<100	<100	<100	<100	<100	<100	<100	<100	<100	52	<100	
4-Nitrophenol	<40	<40	<40	<40	<40	<40	<40	<40	<40	<40	0	<40	
<i>n</i> -Nitroso di- <i>n</i> -propylamine	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	53	<10	
<i>n</i> -Nitrosodiphenylamine	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	68	<10	
Pentachloroanisole	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	73	<20	
Pentachloronitrobenzene	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	88	<30	
Pentachlorophenol	<100	<100	<100	<100	<100	<100	<100	<100	<100	<100	0	<100	
Phenanthrene	<5	<5	31	<5	<5	43	36	<5	<5	<5	71	530	
Phenanthridine	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	74	<10	
Phenol	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	55	200	
Pyrene	46	49	56	<5	<5	53	63	57	44	<5	50	76	
Quinoline	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	58	<10	
2,3,5,6-Tetramethylphenol	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	56	<30	
1,2,4-Trichlorobenzene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	60	<10	
2,4,6-Trichlorophenol	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	15	<20	
2,3,6-Trimethylphthalane	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	62	230	
2,4,6-Trimethylphenol	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	53	<30	
Surrogate samples, in percent recovery													
Nitrobenzene-d5	33	55	20	29	39	48	54	40	60	14	73	31	
2-Fluorobiphenyl	49	62	52	57	53	68	58	55	64	68	61	49	
Terphenyl-d14	52	69	65	62	53	77	60	61	66	68	64	52	
³ Benzo-E pyrene-d12	87	97	75	70	64	114	95	92	108	87	102	74	
Certified concentrations,													
¹ National Institute of Standards and Technology (1990).													
² This surrogate compound is added just before the gel permeation chromatography (GPC) step to monitor for GPC-specific losses (E.T. Furlong, U.S. Geological Survey, written commun., 1993).													

EXPLANATION OF TABLE 12

ABBREVIATIONS AND REPORTING UNITS FOR CHEMICAL CONSTITUENTS AND NOTATIONS
USED IN TABLE

MS	Mississippi
SRM	Standard reference material
rep	Replicate sample
dup	Duplicate sample split in laboratory
RS	Reagent spike sample, in percent recovery
MC	Measured concentration
EC	Expected concentration
STD	Standard deviation
<	Less than
—	No data

Detections shown in bold. Analyzed at the U.S. Geological Survey laboratory, Arvada, Colorado.

Table 12. Concentrations of semivolatile organic compounds in the less than 2-millimeter size fraction of postflood sediment samples in the Mississippi River flood plain, sample set reagent spike, standard reference material, and surrogate samples

Compound	Site number (fig. 1)										SRM			
	MS.1			MS.7			MS.11			MS.13				
	Sam- ple	Rep	MS.3	MS.5	Sam- ple	Rep	MS.9	Sam- ple	Dup	Sam- ple	Rep	RS		
Concentration, in micrograms per kilogram														
Acenaphthene.....	<10	<10	34	<10	<10	<10	<10	<10	<10	<10	<10	67	<10	52
Acenaphthylene.....	<5	<5	48	<5	<5	<5	<5	<5	<5	<5	<5	65	280	115
Acridine.....	<10	<10	74	<10	<10	<10	<10	<10	<10	<10	<10	65	<10	10
Anthracene.....	<5	<5	89	<20	<20	<20	<20	<20	<20	<20	<20	69	410	202
Anthraquinone.....	<20	<20										400	—	42
Azo-benzene.....	<20	<20	20	<20	<20	<20	<20	<20	<20	<20	<20	66	<20	—
Benzof[a]anthracene.....	<10	<10	140	<10	67	63	<10	62	60	<10	<10	78	670	2550
Benzof[b]fluoranthene.....	<10	<10	130	<10	84	<10	<10	76	75	<10	<10	56	760	2780
Benzof[k]fluoranthene.....	<10	<10	100	<10	70	<10	<10	62	62	<10	<10	58	700	2444
Benzof[g,h,i]perylene.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	20	260	2516
Benzof[al]pyrene.....	<10	<10	160	<10	<10	<10	<10	100	100	<10	<10	60	860	670
Benzof[c]quinoline.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	72	<10	130
2,2'-Biquinoline.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	63	<20	—
4-Bromophenyl phenyl ether.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	68	<10	—
Butylbenzylphthalate.....	<10	<10	<10	<10	<10	<10	<10	85	88	<10	<10	76	<10	—
C ₈ -Alkyl-phenol.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	72	<10	—
9H-Carbazole.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	72	190	—
bis(2-Chloroethoxy)methane.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	71	<10	—
bis(2-Chloroisopropyl)ether.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	73	<20	—
4-Chloro-3-methylphenol.....	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	52	<30	—
2-Chloronaphthalene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	65	<10	—
2-Chlorophenol.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	62	<10	—
4-Chlorophenyl phenyl ether.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	67	<10	—
Chrysene.....	<10	<10	180	<10	75	76	<10	69	68	<10	<10	54	80	449
p-Cresol.....	<10	<10	64	<10	<10	<10	<10	<10	<10	<10	<10	10	66	<10
Dibenzof[a,h]anthracene.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	29	<20	—
Dibenzothiophene.....	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	<5	5	69	210
Di-n-butylphthalate.....	60	71	87	65	77	69	72	64	68	77	76	460	—	—
1,2-Dichlorobenzene.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	74	<10	—

Table 12. Concentrations of semivolatile organic compounds in the less than 2-millimeter size fraction of postflood sediment samples in the Mississippi River flood plain, sample set reagent spike, standard reference material, and surrogate samples—Continued

Compound	Site number (fig. 1)										SRM				
	MS.1		MS.3		MS.5		MS.7		MS.9		MS.11		MS.13		
	Sam- ple	Rep	Sam- ple	Rep	Sam- ple	Rep	Sam- ple	Rep	Sam- ple	Rep	Sam- ple	Rep	MC	¹ EC	¹ STD
Concentration, in micrograms per kilogram—Continued															
1,3-Dichlorobenzene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	66	<10	—
1,4-Dichlorobenzene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	68	<10	—
2,4-Dichlorophenol	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	62	<20	—
Diethyl phthalate	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	72	<10	—
1,2-Dimethyl naphthalene	<10	<10	19	<10	<10	<10	<10	<10	<10	<10	<10	<10	66	130	—
1,6-Dimethyl naphthalene	<10	<10	39	<10	<10	<10	<10	<10	<10	<10	<10	<10	66	310	—
2,6-Dimethyl naphthalene	42	39	58	23	50	29	23	25	30	24	66	270	198	23	—
3,5-Dimethylphenol	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	60	<20	—
Dimethyl phthalate	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	66	<10	—
4,6-Dinitro-2-methylphenol	<200	<200	<200	<200	<200	<200	<200	<200	<200	<200	<200	<200	0	<200	—
2,4-Dinitrophenol	<300	<300	<300	<300	<300	<300	<300	<300	<300	<300	<300	<300	0	<300	—
2,4-Dinitrothiophene	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	62	<20	—
2,6-Dinitrotoluene	<50	<50	<50	<50	<50	<50	<50	<50	<50	<50	<50	<50	66	<50	—
Di-n-octyl phthalate	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	61	<20	—
bis(2-Ethyhexyl)phthalate	75	86	220	70	91	80	130	140	74	73	81	2,300	—	—	—
2-Ethyl naphthalene	<10	<10	23	<10	<10	<10	<10	<10	<10	<10	<10	<10	65	120	—
Fluoranthene	54	57	270	45	73	85	43	67	49	53	73	1,000	2,1220	240	—
9H-Fluorene	<10	<10	47	<10	<10	<10	<10	<10	<10	<10	<10	<10	66	240	5
Hexachlorobenzene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	70	<10	—
Hexachlorobutadiene	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	65	<20	—
Hexachlorocyclopentadiene	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	63	<20	—
Hexachloroethane	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	64	<20	—
Indeno[1,2,3-cd]pyrene	<20	<20	69	<20	<20	<20	<20	<20	<20	<20	<20	<20	28	440	2569
Isophorone	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	66	<10	—
Isoquolinoline	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	59	<10	—
1-Methyl-9H-fluorene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	68	<10	—
2-Methylanthracene	<10	<10	83	<10	<10	<10	<10	<10	<10	<10	<10	<10	70	360	7
4,5-Methylenephenanthrene	<20	<20	60	<20	40	41	<20	<20	<20	<20	<20	<20	69	270	—
1-Methylphenanthrene	<10	<10	53	<10	<10	<10	<10	<10	<10	<10	<10	<10	70	280	6
1-Methylpyrene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	72	390	—
Naphthalene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	66	740	1,322
Nitrobenzene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	68	<10	—
2-Nitrophenanthrene	<100	<100	<100	<100	<100	<100	<100	<100	<100	<100	<100	<100	66	<100	—
4-Nitrophenol	<40	<40	<40	<40	<40	<40	<40	<40	<40	<40	<40	<40	0	<40	—
n-Nitrosoodi-n-propylamine.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	54	<10	—

Table 12. Concentrations of semivolatile organic compounds in the less than 2-millimeter size fraction of postflood sediment samples in the Mississippi River flood plain, sample set reagent spike, standard reference material, and surrogate samples—Continued

Compound	Site number (fig. 1)										SRM						
	MS.1					MS.7					MS.13						
	Sam- ple	Rep	MS.3	MS.5	Concentration, in micrograms per kilogram—Continued	Sam- ple	Rep	MS.9	Sam- ple	Dup	MS.11	Sam- ple	Rep	RS	MC	1EC	1STD
<i>n</i> -Nitrosodiphenylamine.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	73	<10	—	—
Pentachloroanisole.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	68	<20	—	—
Pentachloronitrobenzene	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	78	<30	—	—
Pentachlorophenol	<100	<100	<100	<100	<100	<100	<100	<100	<100	<100	<100	<100	<100	0	<100	—	—
Phenanthrene.....	32	<5	160	<5	46	<5	45	44	<5	<5	<5	<5	<5	70	560	2577	59
Phenanthridine	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	71	<10	—	—
Phenol.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	65	210	—	—
Pyrene.....	57	60	220	52	84	87	55	75	71	58	61	74	960	2,1080	200	—	—
Quinoline.....	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	67	<10	—	—
2,3,5,6-Tetramethylphenol	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	60	<30	—	—
1,2,4-Trichlorobenzene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	60	<10	—	—
2,4,6-Trichlorophenol.....	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	60	<20	—	—
2,3,6-Trimethylnaphthalene	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	<10	16	<10	—	—
2,4,6-Trimethylphenol.....	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	<30	63	<30	—	—
Surrogate samples, percent recovery																	
Nitrobenzene-d5.....	9.3	14	23	19	99	16	12	20	25	22	10	45	30	—	—	—	
2-Fluorobiphenyl.....	55	51	52	38	54	55	45	48	46	53	54	55	55	—	—	—	
Terphenyl-d14	60	62	63	45	59	67	64	64	60	64	71	58	64	64	—	—	
Benzo-E-pyrene-d12.....	87	84	93	66	92	96	82	90	91	89	92	65	67	67	—	—	

¹National Institute of Standards and Technology (1990).

²Certified concentrations.

³This surrogate compound is added just before the gel permeation chromatography (GPC) step to monitor for GPC-specific losses (E.T. Furlong, U.S. Geological Survey, written commun., 1993).

Table 13. Tentatively identified compounds that were detected in the gas chromatography/mass spectrometry analysis

[MO, Missouri; MS, Mississippi; —, no established confidence level; >C₂₀, more than 20 carbon molecular chain; 1 (in "Confidence level" column), tentative identification based on an indication of a specific library spectrum; 2 (in "Confidence level" column), confident identification, which means good agreement was observed between the unknown compound and a specific library spectrum. Analyzed at the Enseco Analytical Laboratory, Arvada, Colorado]

Compound	Confidence level	Estimated concentration, in milligrams per kilogram
Missouri River		
MO.1		
Oxygenated hydrocarbon.....	—	0.15
Saturated hydrocarbon: >C ₁₀ -C ₂₀	—	.16
—do—	—	.24
Saturated hydrocarbon: >C ₂₀	—	.26
—do—	—	.31
Saturated hydrocarbon: >C ₂₀	—	.38
—do—	—	.46
—do—	—	.54
—do—	—	.20
Siloxane	—	.29
Saturated hydrocarbon: >C ₂₀	—	.25
—do—	—	.17
Siloxane	—	.40
MO.2		
Saturated hydrocarbon: >C ₂₀	—	0.15
MO.3		
Saturated hydrocarbon: C ₁₀ -C ₂₀	—	0.16
(4aS-cis)-2,3,4,4a,5,6,7,8-octahydro-1,1a,4a,7-tetramethyl-1H-benzocyclo-hepten-7-ol	1	.15
Siloxane	—	.19
Saturated hydrocarbon: >C ₂₀	—	.16
MO.4 (layer 1)		
No compounds detected.....		
MO.4 (layer 2)		
No compounds detected.....		
MO.5		
Saturated hydrocarbon: >C ₂₀	—	0.14
Siloxane	—	.24
—do—	—	.62
—do—	—	2.1
—do—	—	4.1

Table 13. Tentatively identified compounds that were detected in the gas chromatography/mass spectrometry analysis—Continued

[MO, Missouri; MS, Mississippi; —, no established confidence level; >C₂₀, more than 20 carbon molecular chain; 1 (in "Confidence level" column), tentative identification based on an indication of a specific library spectrum; 2 (in "Confidence level" column), confident identification, which means good agreement was observed between the unknown compound and a specific library spectrum. Analyzed at the Enseco Analytical Laboratory, Arvada, Colorado]

Compound	Confidence level	Estimated concentration, in milligrams per kilogram
MO.6		
Saturated hydrocarbon: C ₁₀ -C ₂₀	—	0.15
(Z)-14-Methyl-8-hexadecen-1-ol	1	.16
Cyclododecanol	1	.19
9-Hexadecenoic acid	—	.19
Hexadecanoic acid	—	.65
Siloxane	—	.37
—do—	—	.87
—do—	—	.89
4,8,12,2-Trimethyl-3,7,		
11-tridecatrienenitrile	1	.18
Saturated hydrocarbon: >C ₂₀	—	.23
Siloxane	—	9.1
2-Methyl-1-hexadecanol	1	.22
Saturated hydrocarbon: >C ₂₀	—	.20
Hexadecyl ester tetradecanoic acid	1	.68
Saturated hydrocarbon: >C ₂₀	—	.20
Siloxane	—	.20
Hexadecyl ester hexadecanoic acid	1	.65
MO.7		
Siloxane	—	0.51
3-Ethyl-5-(2-ethylbutyl)-octadecane	1	.14
Siloxane	—	.69
MO.7 (replicate)		
Unknown	—	0.39
Siloxane	—	.32
MO.8		
Saturated hydrocarbon: >C ₂₀	—	0.26
—do—	—	.21
Hexadecanoic acid	1	.34
MO.9		
No compounds detected		
MO.10		
No compounds detected		

Table 13. Tentatively identified compounds that were detected in the gas chromatography/mass spectrometry analysis—Continued

[MO, Missouri; MS, Mississippi; —, no established confidence level; >C₂₀, more than 20 carbon molecular chain; 1 (in "Confidence level" column), tentative identification based on an indication of a specific library spectrum; 2 (in "Confidence level" column), confident identification, which means good agreement was observed between the unknown compound and a specific library spectrum. Analyzed at the Enseco Analytical Laboratory, Aryada, Colorado]

Compound	Confidence level	Estimated concentration, in milligrams per kilogram
MO.11		
Saturated hydrocarbon: >C ₂₀	—	0.14
—do—	—	.14
11-Diene selina-4(14)	1	.22
MO.12		
No compounds detected.....		
Mississippi River		
MS.1		
2-Methyl-1,3-dioxolane-2-propanol	1	6.0
2,4-Dimethyl-2-pentanol	1	.21
Oxygenated hydrocarbon.....	—	.60
Saturated hydrocarbon: >C ₂₀	—	.28
1,2-Dibromo-2-methyl-propane.....	1	.18
Saturated hydrocarbon: >C ₂₀	—	.40
Oxygenated hydrocarbon.....	—	.32
Saturated hydrocarbon: >C ₂₀	—	.82
—do—	—	.37
Oxygenated hydrocarbon.....	—	.24
MS.1 (replicate)		
2-Methyl-1,3-dioxolane-2-propanol	1	6.1
2,4-Dimethyl-2-pentanol	1	.14
Oxygenated hydrocarbon.....	—	.54
—do—	—	.26
—do—	—	.28
Oxygenated hydrocarbon.....	—	.20
Saturated hydrocarbon: >C ₂₀	—	.34
Unsaturated hydrocarbon.....	—	.47
Saturated hydrocarbon: >C ₂₀	—	.59
Oxygenated hydrocarbon.....	—	.26
Saturated hydrocarbon: >C ₂₀	—	.35
MS.2		
2-Methyl-1,3-dioxolane-2-propanol	1	5.1
2,4-Dimethyl-2-pentanol	1	.17
Oxygenated hydrocarbon.....	—	.49
—do—	—	.44
Saturated hydrocarbon: >C ₂₀	—	.16

Table 13. Tentatively identified compounds that were detected in the gas chromatography/mass spectrometry analysis—Continued

[MO, Missouri; MS, Mississippi; —, no established confidence level; >C₂₀, more than 20 carbon molecular chain; 1 (in "Confidence level" column), tentative identification based on an indication of a specific library spectrum; 2 (in "Confidence level" column), confident identification, which means good agreement was observed between the unknown compound and a specific library spectrum. Analyzed at the Enseco Analytical Laboratory, Arvada, Colorado]

Compound	Confidence level	Estimated concentration, in milligrams per kilogram
MS.2—Continued		
1,2-Dibromo-2-methyl-propane	1	0.16
Saturated hydrocarbon: >C ₂₀	—	.14
—do—	—	.68
Oxygenated hydrocarbon	—	.26
Saturated hydrocarbon: >C ₂₀	—	.22
Oxygenated hydrocarbon	—	.34
Saturated hydrocarbon: >C ₂₀	—	1.3
Oxygenated hydrocarbon	—	.83
Saturated hydrocarbon: >C ₂₀	—	.86
Oxygenated hydrocarbon	—	.64
Unsaturated hydrocarbon	—	.88
Oxygenated hydrocarbon	—	.85
—do—	—	.71
MS.3		
2-Methyl-1,3-dioxolane-2-propanol.....	1	5.1
Oxygenated hydrocarbon	—	.58
—do—	—	.69
Saturated hydrocarbon: >C ₂₀	—	.84
—do—	—	1.1
Unsaturated hydrocarbon	—	.56
Saturated hydrocarbon: >C ₂₀	—	1.5
Unsaturated hydrocarbon	—	1.4
Saturated hydrocarbon: >C ₂₀	—	1.1
Oxygenated hydrocarbon	—	.34
Oxygenated hydrocarbon	—	.34
—do—	—	.43
—do—	—	.34
—do—	—	.40
—do—	—	.49
Saturated hydrocarbon: >C ₂₀	—	.85
Oxygenated hydrocarbon	—	.73
Unsaturated hydrocarbon	—	.39
Oxygenated hydrocarbon	—	1.3

Table 13. Tentatively identified compounds that were detected in the gas chromatography/mass spectrometry/analysis—Continued

[MO, Missouri; MS, Mississippi; —, no established confidence level; >C₂₀, more than 20 carbon molecular chain; 1 (in "Confidence level" column), tentative identification based on an indication of a specific library spectrum; 2 (in "Confidence level" column), confident identification, which means good agreement was observed between the unknown compound and a specific library spectrum. Analyzed at the Enseco Analytical Laboratory, Arvada, Colorado]

Compound	Confidence level	Estimated concentration, in milligrams per kilogram
MS.4		
Oxygenated hydrocarbon	—	1.2
—do—	—	.61
—do—	—	.18
—do—	—	1.4
Saturated hydrocarbon: >C ₂₀	—	.49
Saturated hydrocarbon: >C ₂₀	—	.86
—do—	—	.68
—do—	—	1.7
Unsaturated hydrocarbon	—	.58
Oxygenated hydrocarbon	—	.57
Saturated hydrocarbon: >C ₂₀	—	1.1
Oxygenated hydrocarbon	—	.53
—do—	—	.70
—do—	—	1.1
—do—	—	.40
Oxygenated hydrocarbon	—	1.1
—do—	—	1.2
—do—	—	.38
—do—	—	1.5
—do—	—	.40
MS.5		
2-Methyl-1,3-dioxolane-2-propanol	1	5.0
Oxygenated hydrocarbon	—	1.3
—do—	—	.64
—do—	—	1.6
Saturated hydrocarbon: >C ₂₀	—	.52
Diisooctyl ester-1,2-benzenedicarboxlic acid.....	1	.41
Saturated hydrocarbon: >C ₂₀	—	.41
—do—	—	.75
Oxygenated hydrocarbon	—	.46
Saturated hydrocarbon: >C ₂₀	—	.56
Unsaturated hydrocarbon	—	1.8
Saturated hydrocarbon: >C ₂₀	—	1.1
Saturated hydrocarbon: >C ₂₀	—	.82
Oxygenated hydrocarbon	—	.75
—do—	—	.53
Saturated hydrocarbon: >C ₂₀	—	.49
Oxygenated hydrocarbon	—	.88

Table 13. Tentatively identified compounds that were detected in the gas chromatography/mass spectrometry analysis—Continued

[MO, Missouri; MS, Mississippi; —, no established confidence level; >C₂₀, more than 20 carbon molecular chain; 1 (in "Confidence level" column), tentative identification based on an indication of a specific library spectrum; 2 (in "Confidence level" column), confident identification, which means good agreement was observed between the unknown compound and a specific library spectrum. Analyzed at the Enseco Analytical Laboratory, Arvada, Colorado]

Compound	Confidence level	Estimated concentration, in milligrams per kilogram
MS.6		
2-Methyl-1,3-dioxolane-2-propanol.....	1	3.9
Saturated hydrocarbon: C ₁₀ -C ₂₀	—	.68
Oxygenated hydrocarbon	—	.77
—do—	—	1.6
—do—	—	1.5
Oxygenated hydrocarbon	—	.65
—do—	—	1.7
—do—	—	.75
—do—	—	1.2
—do—	—	1.1
Oxygenated hydrocarbon	—	2.8
—do—	—	1.1
—do—	—	1.1
—do—	—	.75
—do—	—	1.0
Oxygenated hydrocarbon	—	1.2
—do—	—	.88
—do—	—	.65
—do—	—	1.8
—do—	—	.97
MS.7		
5-(Acetoxy)-2-pentanone	1	4.1
Oxygenated hydrocarbon	—	.39
Saturated hydrocarbon: >C ₂₀	2	.64
Oxygenated hydrocarbon	—	.56
Saturated hydrocarbon: >C ₂₀	—	.93
Oxygenated hydrocarbon	—	.22
Saturated hydrocarbon: >C ₂₀	—	.27
1,1'-Thiobis-dodecane.....	1	.27
Saturated hydrocarbon: >C ₂₀	—	.92
Cyclic hydrocarbon	—	.21
Unsaturated hydrocarbon	—	.46
Saturated hydrocarbon: >C ₂₀	—	.71
Oxygenated hydrocarbon	—	.25
—do—	—	.61
Saturated hydrocarbon: >C ₂₀	—	.52

Table 13. Tentatively identified compounds that were detected in the gas chromatography/mass spectrometry analysis—Continued

[MO, Missouri; MS, Mississippi; —, no established confidence level; >C₂₀, more than 20 carbon molecular chain; 1 (in "Confidence level" column), tentative identification based on an indication of a specific library spectrum; 2 (in "Confidence level" column), confident identification, which means good agreement was observed between the unknown compound and a specific library spectrum. Analyzed at the Enseco Analytical Laboratory, Arvada, Colorado]

Compound	Confidence level	Estimated concentration, in milligrams per kilogram
MS.7—Continued		
Unsaturated hydrocarbon.....	—	0.67
Oxygenated hydrocarbon.....	—	.25
—do—.....	—	.64
—do—.....	—	.28
MS.7 (replicate)		
2-Methyl-1,3-dioxolane-2-propanol	1	3.5
Oxygenated hydrocarbon.....	—	.20
—do—.....	—	.27
—do—.....	—	.34
—do—.....	—	1.3
Saturated hydrocarbon: >C ₂₀	—	.32
—do—.....	—	.53
—do—.....	—	.21
Oxygenated hydrocarbon.....	—	.21
Saturated hydrocarbon: >C ₂₀	—	.84
Oxygenated hydrocarbon.....	—	.23
Saturated hydrocarbon: >C ₂₀	—	.81
Oxygenated hydrocarbon.....	—	.41
—do—.....	—	.59
Saturated hydrocarbon: >C ₂₀	—	.33
Oxygenated hydrocarbon.....	—	.68
—do—.....	—	.28
—do—.....	—	.66
—do—.....	—	.21
MS.8		
2-Methyl-1,3-dioxolane-2-propanol	1	6.2
Saturated hydrocarbon: C ₁₀ -C ₂₀	—	.48
Oxygenated hydrocarbon.....	—	.57
Unsaturated hydrocarbon.....	—	.25
Oxygenated hydrocarbon.....	—	.31
Oxygenated hydrocarbon.....	—	1.6
—do—.....	—	1.1
—do—.....	—	.59
—do—.....	—	1.5
—do—.....	—	.33

Table 13. Tentatively identified compounds that were detected in the gas chromatograph//mass spectrometry analysis—Continued

[MO, Missouri; MS, Mississippi; —, no established confidence level; >C₂₀, more than 20 carbon molecular chain; 1 (in "Confidence level" column), tentative identification based on an indication of a specific library spectrum; 2 (in "Confidence level" column), confident identification, which means good agreement was observed between the unknown compound and a specific library spectrum. Analyzed at the Enseco Analytical Laboratory, Arvada, Colorado]

Compound	Confidence level	Estimated concentration, in milligrams per kilogram
MS.8—Continued		
Saturated hydrocarbon: >C ₂₀	—	0.27
Oxygenated hydrocarbon	—	.35
—do—.....	—	.36
—do—.....	—	.64
—do—.....	—	.53
Unsaturated hydrocarbon	—	.33
Oxygenated hydrocarbon	—	.88
—do—.....	—	.78
—do—.....	—	.32
MS.9		
Ethyl ester, acetic acid.....	1	0.14
2-Methyl-1,3-dioxolane-2-propanol.....	1	4.4
2,4-Dimethyl-2-pentanol	1	.16
Saturated hydrocarbon: C ₁₀ -C ₂₀	—	.20
Oxygenated hydrocarbon	—	.20
Oxygenated hydrocarbon	—	.51
—do—.....	—	.18
Saturated hydrocarbon: >C ₂₀	—	.15
—do—.....	—	.22
Oxygenated hydrocarbon	—	.24
Saturated hydrocarbon: >C ₂₀	—	.59
—do—.....	—	.34
Aromatic hydrocarbon.....	—	.60
MS.10		
4-(Acetoxy)-2-butanone	1	0.16
2-Methyl-1,3-dioxolane-2-propanol.....	1	5.5
2,4-Dimethyl-2-pentanol	1	.20
Saturated hydrocarbon: C ₁₀ -C ₂₀	—	.32
Oxygenated hydrocarbon	—	.24
Oxygenated hydrocarbon	—	.32
Saturated hydrocarbon: >C ₂₀	—	.16
Unsaturated hydrocarbon	—	.43
Saturated hydrocarbon: >C ₂₀	—	.32
—do—.....	—	.19
Saturated hydrocarbon: >C ₂₀	—	.45
Oxygenated hydrocarbon	—	.24

Table 13. Tentatively identified compounds that were detected in the gas chromatography/mass spectrometry analysis—Continued

[MO, Missouri; MS, Mississippi; —, no established confidence level; >C₂₀, more than 20 carbon molecular chain; 1 (in "Confidence level" column), tentative identification based on an indication of a specific library spectrum; 2 (in "Confidence level" column), confident identification, which means good agreement was observed between the unknown compound and a specific library spectrum. Analyzed at the Enseco Analytical Laboratory, Arvada, Colorado]

Compound	Confidence level	Estimated concentration, in milligrams per kilogram
MS.11		
Saturated hydrocarbon: >C ₂₀	—	0.16
(Z,E)-3,7,11-trimethyl-2,6,10-dodecatrien-1-o1	1	.18
Saturated hydrocarbon: >C ₂₀	—	.33
—do—	—	.24
Siloxane	—	.17
Siloxane	—	.34
MS.12		
4-Methyl-, (Z)-2-pentene	1	0.19
Saturated hydrocarbon: C ₁₀ -C ₂₀	—	.23
Siloxane	—	.33
9-Hexadecenoic acid.....	1	1.3
Hexadecanoic acid	1	1.7
1-Methyl-4-(1-methylethyl)-cyclohexanol	1	.14
5-Bromo-2,3-dimethyl-2-pentene	1	.16
[R- [(E,E)]-3,7,11,15-tetramethyl-6,10,14-hexadecatrien-1-o1.....	1	.23
Saturated hydrocarbon: >C ₂₀	—	.25
1-(Ethenyloxy)-octadecane	1	.17
Saturated hydrocarbon: >C ₂₀	—	.28
Siloxane	—	.26
1,2-Dimethyl ester 1,2,4-benzenetricarboxylic acid.....	1	.30
Alloaromadendrene.....	1	.26
MS.13		
2,4-Dimethyl-3-pantanone	1	0.13
Saturated hydrocarbon: C ₁₀ -C ₂₀	—	.16
Siloxane	—	.34
9-Hexadecenoic acid.....	1	1.4
Hexadecanoic acid	1	1.5
[R- [R,R-(E)]]-3,7,11,15-tetramethyl-2-hexadecen-1-o1	1	.14
3,7,11-Trimethyl-2,6,10-dodecatrien-1-o1	1	.22
Saturated hydrocarbon: >C ₂₀	—	.27
—do—	—	.20
[S-(Z)]-3,7,11-trimethyl-1,6,10-dodecatrien-3-o1	1	.38
Saturated hydrocarbon: >C ₂₀	—	.28
(+)-6-methyl-2-(4-methyl-3-cyclohexen-1-yl)-5-hepten-2-o1	1	.30

Table 13. Tentatively identified compounds that were detected in the gas chromatograph/mass spectrometry analysis—Continued

[MO, Missouri; MS, Mississippi; —, no established confidence level; >C₂₀, more than 20 carbon molecular chain; 1 (in "Confidence level" column), tentative identification based on an indication of a specific library spectrum; 2 (in "Confidence level" column), confident identification, which means good agreement was observed between the unknown compound and a specific library spectrum. Analyzed at the Enseco Analytical Laboratory, Arvada, Colorado]

Compound	Confidence level	Estimated concentration, in milligrams per kilogram
MS.13 (replicate)		
Saturated hydrocarbon: >C ₂₀	—	0.19
Octadecanal	1	.16
3,7,11-Trimethyl-2,6,10-dodecatrien-1-ol.....	1	.21
Saturated hydrocarbon: >C ₂₀	—	.39
—do—.....	—	.35
1,1'-ethylenbis-cyclohexane.....	1	.19
5,6-dihydro-5,6-dihydroxy-.beta.-carotene. beta	1	.31
Sclareol.....	1	.51

Selected Series of U.S. Geological Survey Publications

Books and Other Publications

Professional Papers report scientific data and interpretations of lasting scientific interest that cover all facets of USGS investigations and research.

Bulletins contain significant data and interpretations that are of lasting scientific interest but are generally more limited in scope or geographic coverage than Professional Papers.

Water-Supply Papers are comprehensive reports that present significant interpretive results of hydrologic investigations of wide interest to professional geologists, hydrologists, and engineers. The series covers investigations in all phases of hydrology, including hydrogeology, availability of water, quality of water, and use of water.

Circulars are reports of programmatic or scientific information of an ephemeral nature; many present important scientific information of wide popular interest. Circulars are distributed at no cost to the public.

Fact Sheets communicate a wide variety of timely information on USGS programs, projects, and research. They commonly address issues of public interest. Fact Sheets generally are two or four pages long and are distributed at no cost to the public.

Reports in the **Digital Data Series (DDS)** distribute large amounts of data through digital media, including compact disc-read-only memory (CD-ROM). They are high-quality, interpretive publications designed as self-contained packages for viewing and interpreting data and typically contain data sets, software to view the data, and explanatory text.

Water-Resources Investigations Reports are papers of an interpretive nature made available to the public outside the formal USGS publications series. Copies are produced on request (unlike formal USGS publications) and are also available for public inspection at depositories indicated in USGS catalogs.

Open-File Reports can consist of basic data, preliminary reports, and a wide range of scientific documents on USGS investigations. Open-File Reports are designed for fast release and are available for public consultation at depositories.

Maps

Geologic Quadrangle Maps (GQ's) are multicolor geologic maps on topographic bases in 7.5- or 15-minute quadrangle formats (scales mainly 1:24,000 or 1:62,500) showing bedrock, surficial, or engineering geology. Maps generally include brief texts; some maps include structure and columnar sections only.

Geophysical Investigations Maps (GP's) are on topographic or planimetric bases at various scales. They show results of geophysical investigations using gravity, magnetic, seismic, or radioactivity surveys, which provide data on subsurface structures that are of economic or geologic significance.

Miscellaneous Investigations Series Maps or Geologic Investigations Series (I's) are on planimetric or topographic bases at various scales; they present a wide variety of format and subject matter. The series also includes 7.5-minute quadrangle photogeologic maps on planimetric bases and planetary maps.

Information Periodicals

Metal Industry Indicators (MII's) is a free monthly newsletter that analyzes and forecasts the economic health of five metal industries with composite leading and coincident indexes: primary metals, steel, copper, primary and secondary aluminum, and aluminum mill products.

Mineral Industry Surveys (MIS's) are free periodic statistical and economic reports designed to provide timely statistical data on production, distribution, stocks, and consumption of significant mineral commodities. The surveys are issued monthly, quarterly, annually, or at other regular intervals, depending on the need for current data. The MIS's are published by commodity as well as by State. A series of international MIS's is also available.

Published on an annual basis, **Mineral Commodity Summaries** is the earliest Government publication to furnish estimates covering nonfuel mineral industry data. Data sheets contain information on the domestic industry structure, Government programs, tariffs, and 5-year salient statistics for more than 90 individual minerals and materials.

The Minerals Yearbook discusses the performance of the worldwide minerals and materials industry during a calendar year, and it provides background information to assist in interpreting that performance. The Minerals Yearbook consists of three volumes. Volume I, Metals and Minerals, contains chapters about virtually all metallic and industrial mineral commodities important to the U.S. economy. Volume II, Area Reports: Domestic, contains a chapter on the minerals industry of each of the 50 States and Puerto Rico and the Administered Islands. Volume III, Area Reports: International, is published as four separate reports. These reports collectively contain the latest available mineral data on more than 190 foreign countries and discuss the importance of minerals to the economies of these nations and the United States.

Permanent Catalogs

"**Publications of the U.S. Geological Survey, 1879-1961**" and "**Publications of the U.S. Geological Survey, 1962-1970**" are available in paperback book form and as a set of microfiche.

"**Publications of the U.S. Geological Survey, 1971-1981**" is available in paperback book form (two volumes, publications listing and index) and as a set of microfiche.

Annual supplements for 1982, 1983, 1984, 1985, 1986, and subsequent years are available in paperback book form.



Printed on recycled paper

ISBN 0-607-88279-4



9 780607 882797